

Product datasheet for MR220367

Arfgap1 (NM_001177708) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Arfgap1 (NM_001177708) Mouse Tagged ORF Clone

Tag: Myc-DDK
Symbol: Arfgap1

Synonyms: Al115377; Arf1gap

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

ORF Nucleotide >MR220367 representing NM_001177708
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Sequence: >MR220367 representing NM_001177708

Red=Cloning site Green=Tags(s)


MASPRTRKVLKEVRAQDENNVCFECGAFNPQWVSVTYGIWICLECSGRHRGLGVHLSFVRSVTMDKWKDI ELEKMKAGGNAKFREFLETQDDYEPSWSLQDKYSSRAAALFRDKVATLAEGKEWSLESSPAQNWTPPQPK TLQFTAHRASGQPQSAAASGDKAFEDWLNDDLGSYQGAQENRYVGFGNTVPPQKREDDFLNNAMSSLYSG WSSFTTGASKFASAAKEGATKFGSQASQKASELGHSPE


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_001177708

ORF Size: 744 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at customercom or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Arfgap1 (NM_001177708) Mouse Tagged ORF Clone - MR220367

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq Size: 2448 bp
RefSeq ORF: 1185 bp
Locus ID: 228998
UniProt ID: Q9EPI9

Cytogenetics: 2 103.53 cM

MW: 43.6 kDa

Gene Summary: GTPase-activating protein (GAP) for the ADP ribosylation factor 1 (ARF1). Involved in


membrane trafficking and /or vesicle transport. Promotes hydrolysis of the ARF1-bound GTP and thus, is required for the dissociation of coat proteins from Golgi-derived membranes and vesicles, a prerequisite for vesicle's fusion with target compartment. Probably regulates ARF1-mediated transport via its interaction with the KDELR proteins and TMED2. Overexpression induces the redistribution of the entire Golgi complex to the endoplasmic reticulum, as when

ARF1 is deactivated. Its activity is stimulated by phosphoinosides and inhibited by

phosphatidylcholine (By similarity).[UniProtKB/Swiss-Prot Function]



Product images:



Circular map for MR220367