

Product datasheet for MR220508

Babam2 (NM_181281) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Babam2 (NM_181281) Mouse Tagged ORF Clone
Tag: Myc-DDK
Symbol: Babam2
Synonyms: 6030405P19Rik; AI429776; B830038C02Rik
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
ORF Nucleotide Sequence: >MR220508 representing NM_181281
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCCGCGATCGCC

ATGTGTGCCTGTAGGGACATAATTTTCAATGCTCAGTACCCAGAGCTGCCTCCTGATTTTCATCTTTGGAG
 AGGATGCTGAGTTTCTGCCAGACCCCTCTGCGCTGCACAACCTTGCTCCTGGAACCTTCAAACCTGA
 GTGCCTGTTGCTCGTGGTGAAGGAGCTGGTGCAGCAGTACCACAGTTCCAGTGCAGCCGCTCCGTGAG
 AGCTCGCGCCTCATGTTTGAGTACCAGACGCTCCTGGAAGAGCCTCAGTATGGAGAGAACATGGAATTT
 ATGCTGGAAGAAAAACAACCTGGACTGGTGAATTTTCAGCTCGTTTTCTATTGAAGTTACCAAGTAGATT
 CAGCAACATCCACATACCTTCTCAAGGATGTAATGAAGACCTGGAGAAGATGTGGCCCTTCTTTCT
 GTCAGTTTTGAGGATACTGAAGTACCCAGGTGTACCCCAAGTTGTACTTGTACCCCGAATTGAGCATG
 CACTCGGAGGCTCCTCTGCTCTTACATCCCTGCTTCCCGGAGGAGGATGTCTCATTGATTATGTGCC
 TCAAGTGTGCCACCTGCTACCAACAAGGTACAGTATGTGATTCAAGGCTATCACAAAAGAAGAGAGTAC
 ATCGCAGCTTTCCTCAGTCACTTTGGCACAGGTGTCGTGGAATATGATGCAGAAGGCTTCACAAAACCTA
 CTCTGCTGCTGATGTGGAAGACTTTTGTCTTGTCCACATTGACCTGCCCTGTTTTCCCTCGAGA
 TCAGCCTACGCTCACATTTTCAGTCACTATCACTTTACCAACAGTGGACAGCTTTACTCCCAAGCACAA
 AAAAATACCCATACAGCCCGAGATGGGATGGAATGAAATGGCCAAGAGAGCAAAGGCTTATTTCAAAA
 CCTTTGCTCCCTCAGTTCCAGGAGCCGCATTTGCCAATGGAAAGCTC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >MR220508 representing NM_181281
 Red=Cloning site Green=Tags(s)

MCACRDIIFNAQYPELPPDFIFGEDAEFLPDPSSALHNLASWNPSNPECLLLVVKELVQQYHQFCGRRLRE
 SSRLMFEYQTLLEEPQYGENMEIYAGKKNWTFGEFSARFLKLPVDFSNIPTYLLKDVNEDPGEDVALLS
 VSFEDTEATQVYPKLYLSPRIEHALGGSSALHIPAFPGGGLIDYVYPQVCHLLTNKVQYVIQGYHKRREY
 IAAFLSHFGTGVVEYDAEGFTKL TLLL MWKDFCF LVHIDLPLFFPRDQPTLTFQSVYHFTNSGQLYSQAQ
 KNYPYSPRW DGNEMAKRAKAYFKTFVPQFQEAFAFANGKL


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:



Plasmid Map:



ACCN: NM_181281

ORF Size: 957 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_181281.1 , NP_851798.1
RefSeq Size:	1143 bp
RefSeq ORF:	960 bp
Locus ID:	107976
UniProt ID:	Q8K3W0
Cytogenetics:	5 B1
MW:	37 kDa
Gene Summary:	Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). The BRCA1-A complex also possesses deubiquitinase activity that specifically removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX. In the BRCA1-A complex, it acts as an adapter that bridges the interaction between BABAM1/NBA1 and the rest of the complex, thereby being required for the complex integrity and modulating the E3 ubiquitin ligase activity of the BRCA1-BARD1 heterodimer. Probably also plays a role as a component of the BRISC complex, a multiprotein complex that specifically cleaves 'Lys-63'-linked ubiquitin (By similarity). May regulate TNF-alpha signaling through its interactions with TNFRSF1A.[UniProtKB/Swiss-Prot Function]