

Product datasheet for **MR220609**

Exosc6 (NM_028274) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Exosc6 (NM_028274) Mouse Tagged ORF Clone
Tag: Myc-DDK
Symbol: Exosc6
Synonyms: 2610510N21Rik; C76919; Mtr3
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >MR220609 representing NM_028274
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCCCGATCGCC

ATGCCGGGAGACCATCGCCGCATCCGCGGGCCGGAGGAGTCTCAGCCGCCGAGCTGTACGCGGCCGAAG
 ACGACGAGACGCCCGTCTCGGACCCGACGCGGCTCCGCCGGTGTACGCGCGCGCCGGGCTGCTGAG
 CCAGGCCAAGGGCTCCGCCTACCTGGAGGCGGGCGGACCAAGGTGCTGTGCGCCGTGCCGGTCCGCGG
 CAGGCCGAGGGCGGCGAGCGCGGACGCGCCCTCGGGAGCGGGCGGGAAGCTCCCGCTGCCCTCCGGG
 GTCGCTGCTCTGCGATTCGCGCGCTCCCTTCTCCGGTCCCGGCGTCCGCGCCCCAGGGCGGCGG
 CGGGGAGGATCGCGAGCTGGGACTGGCGCTACAGGAGGCTCTGGAGCCCGCCGTGCGCCTGGGGCGCTAC
 CCGCGTGCTCAGCTCGAGGTGTCGGCGTCTGCTGGAGGACGGCGGCTGTGCGTGGCCGCGGCGCTCA
 CCGCGGCCGCGCTCGCCCTGGCCGACGCGGGAGTGGAGATGTACGATCTGGTGGTGGGCTGCGGCCTGAG
 CCTCACCCGGGACCCCTCCCGACCTGGCTGCTGGACCCACGCGGCTGGAGGAGGAGCACTCCGCGGCC
 GGTCTCACGGTGGCCCTCATGCCTGTGCTCAATCAGGTGGCCGGGTTACTGGGCAGTGGGAAGGCGGGC
 AGACTGAGAGTTGACCGATGCTGTGCGCCTGGGCTGGAGGGCTGCCAGCGCCTCTACCCAGTGTGCA
 GCAGTGTGGTGGCGGGCTGCCCGCGGAGGGGTGCCCGCCCAACC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >MR220609 representing NM_028274
Red=Cloning site Green=Tags(s)

MPGDHRRIRGPEESQPPQLYAAEDDETPAARDPTRLRPVYARAGLLSQAKGSAYLEAGGTKVLCAVSGPR
 QAEGGERGSGPAGAGGEAPAALRGRLLCDFRRAPFSGRRRRAPQGGGGEDRELGLALQEALPAVRLGRL
 PRAQLEVSALLLEDGGCALAAAL TAAALALADAGVEMYDLVVGGLSLTPGPSPTWLLDPTRL EEEHSAA
 GLTVALMPVLNQVAGLLGSGEGGQTESWTDVAVRLGLEGCQRLYPVLQQCLVRAARRRRGAAAPP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mm9046_a10.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_028274

ORF Size: 819 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_028274.4 , NP_082550.1
RefSeq Size:	1346 bp
RefSeq ORF:	822 bp
Locus ID:	72544
UniProt ID:	Q8BTW3
Cytogenetics:	8 E1
MW:	28.8 kDa
Gene Summary:	<p>Non-catalytic component of the RNA exosome complex which has 3'->5' exoribonuclease activity and participates in a multitude of cellular RNA processing and degradation events. In the nucleus, the RNA exosome complex is involved in proper maturation of stable RNA species such as rRNA, snRNA and snoRNA, in the elimination of RNA processing by-products and non-coding 'pervasive' transcripts, such as antisense RNA species and promoter-upstream transcripts (PROMPTs), and of mRNAs with processing defects, thereby limiting or excluding their export to the cytoplasm. The RNA exosome may be involved in Ig class switch recombination (CSR) and/or Ig variable region somatic hypermutation (SHM) by targeting AICDA deamination activity to transcribed dsDNA substrates. In the cytoplasm, the RNA exosome complex is involved in general mRNA turnover and specifically degrades inherently unstable mRNAs containing AU-rich elements (AREs) within their 3' untranslated regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs. It seems to be involved in degradation of histone mRNA. The catalytic inactive RNA exosome core complex of 9 subunits (Exo-9) is proposed to play a pivotal role in the binding and presentation of RNA for ribonucleolysis, and to serve as a scaffold for the association with catalytic subunits and accessory proteins or complexes (By similarity).[UniProtKB/Swiss-Prot Function]</p>

Product images:



Circular map for MR220609