

## Product datasheet for MR220692L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Ube2o (NM\_173755) Mouse Tagged ORF Clone Lentiviral Particle**

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** Ube2o (NM\_173755) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Ube2d

**Synonyms:** 9630022H21; B230113M03Rik; E2-230K; mKIAA1734

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 173755

ORF Size: 3864 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR220692).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 173755.3, NP 776116.2

RefSeq Size: 5192 bp
RefSeq ORF: 3867 bp
Locus ID: 217342
UniProt ID: Q6ZPJ3
Cytogenetics: 11 E2





## **Gene Summary:**

E2/E3 hybrid ubiquitin-protein ligase that displays both E2 and E3 ligase activities and mediates monoubiquitination of target proteins. Negatively regulates TRAF6-mediated NF-kappa-B activation independently of its E2 activity. Acts as a positive regulator of BMP7 signaling by mediating monoubiquitination of SMAD6, thereby regulating adipogenesis. Mediates monoubiquitination at different sites of the nuclear localization signal (NLS) of BAP1, leading to cytoplasmic retention of BAP1. Also able to monoubiquitinate the NLS of other chromatin-associated proteins, such as INO80 and CXXC1, affecting their subcellular location. Acts as a regulator of retrograde transport by assisting the TRIM27:MAGEL2 E3 ubiquitin ligase complex to mediate 'Lys-63'-linked ubiquitination of WASHC1, leading to promote endosomal F-actin assembly.[UniProtKB/Swiss-Prot Function]