

Product datasheet for MR221045L4

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

NIrp1a (NM_001004142) Mouse Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Nlrp1a (NM_001004142) Mouse Tagged Lenti ORF Clone

Tag: mGFP Symbol: NIrp1a

Synonyms: CARD7; DEFCAP; Gm14; Gm15; NAC; Nalp1; Nalp1a; Nlrp1; PP1044

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)


ORF Nucleotide The ORF insert of this clone is exactly the same as(MR221045).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_001004142

ORF Size: 3546 bp

Nlrp1a (NM_001004142) Mouse Tagged Lenti ORF Clone - MR221045L4

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 001004142.2</u>, <u>NP 001004142.2</u>

 RefSeq Size:
 4806 bp

 RefSeq ORF:
 3549 bp

 Locus ID:
 195046

 UniProt ID:
 Q2LKU9

Cytogenetics: 11 43.21 cM


Gene Summary: As a potential sensor component of the NLRP1 inflammasome, plays a crucial role in innate

immunity and inflammation. In response to pathogens and other damage-associated signals, initiates the formation of the inflammasome polymeric complex, made of Nlrp1b, CASP1, and possibly PYCARD. Recruitment of proCASP1 to the inflammasome promotes its activation and CASP1-catalyzed IL1B and IL18 maturation and secretion in the extracellular milieu. Activation of NLRP1 inflammasome is also required for HMGB1 secretion. The active cytokines and HMGB1 stimulate inflammatory responses. Inflammasomes can also induce pyroptosis, an inflammatory form of programmed cell death (By similarity) (PubMed:23219391). When activated in the bone marrow, induces the pyroptosis of hematopoietic stem cells and progenitor cells of both myeloid and lymphoid lineages, hence allowing the removal of damaged cells, and the release of IL1B, which induces granulopoiesis (PubMed:23219391).

Binds ATP (By similarity).[UniProtKB/Swiss-Prot Function]



Product images:



Circular map for MR221045L4