

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR221580L4V

Kcnj1 (NM_001168354) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Kcnj1 (NM_001168354) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Kcnj1
Synonyms:	Kir1.1; ROMK; Romk2
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001168354
ORF Size:	1119 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR221580).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	
	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	
RefSeq: RefSeq Size:	varies depending on the nature of the gene.
•	varies depending on the nature of the gene.
RefSeq Size:	varies depending on the nature of the gene. <u>NM_001168354.1</u> 3075 bp



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Kcnj1 (NM_001168354) Mouse Tagged ORF Clone Lentiviral Particle – MR221580L4V
Gene Summary:	In the kidney, probably plays a major role in potassium homeostasis. Inward rectifier
	potassium channels are characterized by a greater tendency to allow potassium to flow into
	the cell rather than out of it. Their voltage dependence is regulated by the concentration of

extracellular potassium; as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the

and can be blocked by external barium (By similarity).[UniProtKB/Swiss-Prot Function]

blockage of outward current by internal magnesium. This channel is activated by internal ATP

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US