

## Product datasheet for MR221618L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Gria4 (NM\_001113180) Mouse Tagged ORF Clone Lentiviral Particle**

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** Gria4 (NM\_001113180) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Gria4

Synonyms: Glu; GluA4; Glur; Glur-4; GluR-D; Glur4; Gluralpha4; spk; spkw1

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_001113180

ORF Size: 2706 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR221618).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** NM 001113180.1, NP 001106651.1

 RefSeq Size:
 5458 bp

 RefSeq ORF:
 2709 bp

 Locus ID:
 14802

 UniProt ID:
 Q9Z2W8

 Cytogenetics:
 9 2.46 cM







## **Gene Summary:**

Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. These receptors are heteromeric protein complexes composed of multiple subunits, arranged to form ligand-gated ion channels. The classification of glutamate receptors is based on their activation by different pharmacologic agonists. The subunit encoded by this gene belongs to a family of AMPA (alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate)-sensitive glutamate receptors, and is subject to RNA editing (AGA->GGA; R->G). Alternative splicing of this gene results in transcript variants encoding different isoforms, which may vary in their signal transduction properties. [provided by RefSeq, Jul 2008]