

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR221796L3V

Magt1 (NM_001190409) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Magt1 (NM_001190409) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Magt1
Synonyms:	2410001C15Rik; 2610529C04Rik; 2810482l07Rik; IAG2; IAP
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001190409
ORF Size:	1104 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR221796).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001190409.1, NP 001177338.1</u>
RefSeq Size:	4524 bp
RefSeq ORF:	1107 bp
Locus ID:	67075
UniProt ID:	<u>Q9CQY5</u>
Cytogenetics:	X D



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Acts as accessory component of the N-oligosaccharyl transferase (OST) complex which
catalyzes the transfer of a high mannose oligosaccharide from a lipid-linked oligosaccharide
donor to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent
polypeptide chains. Involved in N-glycosylation of STT3B-dependent substrates. Specifically
required for the glycosylation of a subset of acceptor sites that are near cysteine residues; in
this function seems to act redundantly with TUSC3. In its oxidized form proposed to form
transient mixed disulfides with a glycoprotein substrate to facilitate access of STT3B to the
unmodified acceptor site. Has also oxidoreductase-independent functions in the STT3B-
containing OST complex possibly involving substrate recognition.[UniProtKB/Swiss-Prot
Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US