

Product datasheet for MR221850L4

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

Rockville, MD 20850, US

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

EU: info-de@origene.com CN: techsupport@origene.cn

Spn (NM_001037810) Mouse Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Spn (NM_001037810) Mouse Tagged Lenti ORF Clone

Tag: mGFP Symbol: Spn

Synonyms: A630014B01Rik; Cd43; Galgp; Ly-48; Ly48

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(MR221850).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_001037810

ORF Size: 1185 bp



Spn (NM_001037810) Mouse Tagged Lenti ORF Clone - MR221850L4

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001037810.1</u>, <u>NP 001032899.1</u>

 RefSeq Size:
 1475 bp

 RefSeq ORF:
 1188 bp

 Locus ID:
 20737

 UniProt ID:
 P15702


Cytogenetics: 7 69.35 cM

Gene Summary:

Predominant cell surface sialoprotein of leukocytes which regulates multiple T-cell functions, including T-cell activation, proliferation, differentiation, trafficking and migration. Positively regulates T-cell trafficking to lymph-nodes via its association with ERM proteins (EZR, RDX and MSN) (PubMed:17638845, PubMed:21289089, PubMed:11728336). Negatively regulates Th2 cell differentiation and predisposes the differentiation of T-cells towards a Th1 lineage commitment (PubMed:18490738). Promotes the expression of IFN-gamma by T-cells during T-cell receptor (TCR) activation of naive cells and induces the expression of IFN-gamma by CD4(+) T-cells and to a lesser extent by CD8(+) T-cells. Plays a role in preparing T-cells for cytokine sensing and differentiation into effector cells by inducing the expression of cytokine receptors IFNGR and IL4R, promoting IFNGR and IL4R signaling and by mediating the clustering of IFNGR with TCR (By similarity). Acts as a major E-selectin ligand responsible for Th17 cell rolling on activated vasculature and recruitment during inflammation. Mediates Th17 cells, but not Th1 cells, adhesion to E-selectin (PubMed:26700769). Acts as a T-cell counter-receptor for SIGLEC1 (PubMed:11238599).[UniProtKB/Swiss-Prot Function]



Product images:



Circular map for MR221850L4