

Product datasheet for **MR221858L3V**

Ipo7 (NM_181517) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Ipo7 (NM_181517) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Ipo7
Synonyms:	A330055O14Rik; C330016G14; Imp7; Ranbp7
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_181517
ORF Size:	3114 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR221858).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_181517.3
RefSeq Size:	3117 bp
RefSeq ORF:	3117 bp
Locus ID:	233726
UniProt ID:	Q9EPL8
Cytogenetics:	7 E3



[View online »](#)

Gene Summary:

Functions in nuclear protein import, either by acting as autonomous nuclear transport receptor or as an adapter-like protein in association with the importin-beta subunit KPNB1. Acting autonomously is thought to serve itself as receptor for nuclear localization signals (NLS) and to promote translocation of import substrates through the nuclear pore complex (NPC) by an energy requiring, Ran-dependent mechanism. At the nucleoplasmic side of the NPC, Ran binds to importin, the importin/substrate complex dissociates and importin is re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran. Mediates autonomously the nuclear import of ribosomal proteins RPL23A, RPS7 and RPL5. Binds to a beta-like import receptor binding (BIB) domain of RPL23A. In association with KPNB1 mediates the nuclear import of H1 histone and the Ran-binding site of IPO7 is not required but synergizes with that of KPNB1 in importin/substrate complex dissociation (By similarity). In vitro, mediates nuclear import of H2A, H2B, H3 and H4 histones.[UniProtKB/Swiss-Prot Function]