

## Product datasheet for **MR221926L3V**

### Syt7 (NM\_018801) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Syt7 (NM_018801) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Syt7
Synonyms:	AI851541; B230112P13Rik
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_018801
ORF Size:	1209 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR221926).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_018801.3</a> , <a href="#">NP_061271.1</a>
RefSeq Size:	6342 bp
RefSeq ORF:	1212 bp
Locus ID:	54525
UniProt ID:	<a href="#">Q9R0N7</a>
Cytogenetics:	19 A



[View online »](#)

**Gene Summary:**

Ca(2+) sensor involved in Ca(2+)-dependent exocytosis of secretory and synaptic vesicles through Ca(2+) and phospholipid binding to the C2 domain. Ca(2+) induces binding of the C2-domains to phospholipid membranes and to assembled SNARE-complexes; both actions contribute to triggering exocytosis. SYT7 binds Ca(2+) with high affinity and slow kinetics compared to other synaptotagmins (PubMed:26738595). Involved in Ca(2+)-triggered lysosomal exocytosis, a major component of the plasma membrane repair (By similarity). Ca(2+)-regulated delivery of lysosomal membranes to the cell surface is also involved in the phagocytic uptake of particles by macrophages (PubMed:16982801, PubMed:21041449). Ca(2+)-triggered lysosomal exocytosis also plays a role in bone remodeling by regulating secretory pathways in osteoclasts and osteoblasts (PubMed:18539119). Involved in cholesterol transport from lysosome to peroxisome by promoting membrane contacts between lysosomes and peroxisomes: probably acts by promoting vesicle fusion by binding phosphatidylinositol-4,5-bisphosphate on peroxisomal membranes (PubMed:25860611). Acts as a key mediator of synaptic facilitation, a process also named short-term synaptic potentiation: synaptic facilitation takes place at synapses with a low initial release probability and is caused by influx of Ca(2+) into the axon terminal after spike generation, increasing the release probability of neurotransmitters (PubMed:24569478, PubMed:26738595). Probably mediates synaptic facilitation by directly increasing the probability of release (PubMed:26738595). May also contribute to synaptic facilitation by regulating synaptic vesicle replenishment, a process required to ensure that synaptic vesicles are ready for the arrival of the next action potential: SYT7 is required for synaptic vesicle replenishment by acting as a sensor for Ca(2+) and by forming a complex with calmodulin (PubMed:24569478). Also acts as a regulator of Ca(2+)-dependent insulin and glucagon secretion in beta-cells (PubMed:18308938, PubMed:19171650). Triggers exocytosis by promoting fusion pore opening and fusion pore expansion in chromaffin cells (PubMed:20956309). Also regulates the secretion of some non-synaptic secretory granules of specialized cells (By similarity). [UniProtKB/Swiss-Prot Function]