

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR222118L3V

Glra1 (NM_020492) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Glra1 (NM_020492) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Glra1
Synonyms:	nmf11; oscillator; ot; spasmodic; spd
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_020492
ORF Size:	1350 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR222118).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 020492.3</u>
RefSeq Size:	2389 bp
RefSeq ORF:	1350 bp
Locus ID:	
Locus ID.	14654



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Glycine receptors are ligand-gated chloride channels. Channel opening is triggered by
extracellular glycine (PubMed:16672662, PubMed:17114051, PubMed:24801766). Channel
opening is also triggered by taurine and beta-alanine (By similarity). Channel characteristics
depend on the subunit composition; heteropentameric channels are activated by lower
glycine levels and display faster desensitization (By similarity). Plays an important role in the
down-regulation of neuronal excitability (PubMed:9145798). Contributes to the generation of
inhibitory postsynaptic currents (PubMed:16672662, PubMed:17114051, PubMed:24801766).
Channel activity is potentiated by ethanol. Potentiation of channel activity by intoxicating
levels of ethanol contribute to the sedative effects of ethanol (PubMed:24801766).
[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US