

Product datasheet for MR222167

MgII (NM_001166251) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
 Product Name: MgII (NM_001166251) Mouse Tagged ORF Clone
 Tag: Myc-DDK
 Symbol: MgII
 Synonyms: AA589436; Ma; MagI; Mgl
 Mammalian Cell Selection: Neomycin
 Vector: pCMV6-Entry (PS100001)
 E. coli Selection: Kanamycin (25 ug/mL)
 Restriction Sites: SgfI-MluI
 Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF


ACCN: NM_001166251
 ORF Size: 957 bp



[View online >](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001166251.1 , NP_001159723.1
RefSeq Size:	3991 bp
RefSeq ORF:	960 bp
Locus ID:	23945
UniProt ID:	O35678
Cytogenetics:	6 D1
MW:	35.7 kDa
Gene Summary:	This gene encodes a monoglyceride lipase, which catalyzes the hydrolysis of monoglycerides into fatty acids and glycerol. This enzyme is also thought to hydrolyze the endocannabinoid 2-arachidonoylglycerol. Alternatively spliced transcript variants have been described. [provided by RefSeq, Oct 2009]

Product images:



Circular map for MR222167