

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for MR222203L4V

## Tpx2 (NM\_001141975) Mouse Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Tpx2 (NM_001141975) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Tpx2
Synonyms:	2610005B21Rik; DIL2; p100; REPP86
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001141975
ORF Size:	2238 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR222203).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001141975.1, NP 001135447.1</u>
RefSeq Size:	4161 bp
RefSeq ORF:	2238 bp
Locus ID:	72119
UniProt ID:	A2APB8
Cytogenetics:	2 H1
cytogenetics.	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Tpx2 (NM_001141975) Mouse Tagged ORF Clone Lentiviral Particle – MR222203L4V
Gene Summary:	Spindle assembly factor required for normal assembly of mitotic spindles. Required for normal assembly of microtubules during apoptosis. Required for chromatin and/or kinetochore dependent microtubule nucleation. Mediates AURKA localization to spindle
	microtubules. Activates AURKA by promoting its autophosphorylation at 'Thr-288' and protects this residue against dephosphorylation. TPX2 is inactivated upon binding to
	importin-alpha. At the onset of mitosis, GOLGA2 interacts with importin-alpha, liberating TPX2
	from importin-alpha, allowing TPX2 to activates AURKA kinase and stimulates local

microtubule nucleation.[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US