

Product datasheet for MR22236L3

Kat2a (NM_020004) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Kat2a (NM_020004) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Kat2a
Synonyms:	1110051E14Rik; AW212720; Gcn5; Gcn5l2; mmGCN5
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR222236).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.

ACCN:	NM_020004
ORF Size:	2487 bp




[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_020004.5 , NP_064388.2
RefSeq Size:	3048 bp
RefSeq ORF:	2493 bp
Locus ID:	14534
UniProt ID:	Q9JHD2
Cytogenetics:	11 63.53 cM

Gene Summary:

Protein lysine acyltransferase that can act both as an acetyltransferase and succinyltransferase, depending on the context (PubMed:28424240). Acts as a histone lysine succinyltransferase: catalyzes succinylation of histone H3 on 'Lys-79' (H3K79succ), with a maximum frequency around the transcription start sites of genes (By similarity). Succinylation of histones gives a specific tag for epigenetic transcription activation (By similarity). Association with the 2-oxoglutarate dehydrogenase complex, which provides succinyl-CoA, is required for histone succinylation (By similarity). In different complexes, functions either as an acetyltransferase (HAT) or as a succinyltransferase: in the SAGA and ATAC complexes, acts as a histone acetyltransferase (By similarity). Has significant histone acetyltransferase activity with core histones, but not with nucleosome core particles (By similarity). Acetylation of histones gives a specific tag for epigenetic transcription activation (PubMed:28424240). Involved in long-term memory consolidation and synaptic plasticity: acts by promoting expression of a hippocampal gene expression network linked to neuroactive receptor signaling (PubMed:25024434). Acts as a positive regulator of T-cell activation: upon TCR stimulation, recruited to the IL2 promoter following interaction with NFATC2 and catalyzes acetylation of histone H3 at Lys-9 (H3K9ac), leading to promote IL2 expression (PubMed:28424240). Also acetylates non-histone proteins, such as CEBPB, PLK4 and TBX5 (PubMed:17301242). Involved in heart and limb development by mediating acetylation of TBX5, acetylation regulating nucleocytoplasmic shuttling of TBX5 (By similarity). Acts as a negative regulator of centrosome amplification by mediating acetylation of PLK4 (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:


Circular map for MR222236L3