

Product datasheet for MR222236L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Kat2a (NM_020004) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Kat2a (NM_020004) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Kat2a

Synonyms: 1110051E14Rik; AW212720; Gcn5; Gcn5l2; mmGCN5

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_020004 **ORF Size:** 2487 bp

ORF Nucleotide

- - -

Sequence:

The ORF insert of this clone is exactly the same as(MR222236).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 020004.5, NP 064388.2</u>

RefSeq Size: 3048 bp
RefSeq ORF: 2493 bp
Locus ID: 14534
UniProt ID: Q9|HD2

Cytogenetics: 11 63.53 cM







Gene Summary:

Protein lysine acyltransferase that can act both as a acetyltransferase and succinyltransferase, depending on the context (PubMed:28424240). Acts as a histone lysine succinyltransferase: catalyzes succinylation of histone H3 on 'Lys-79' (H3K79succ), with a maximum frequency around the transcription start sites of genes (By similarity). Succinylation of histones gives a specific tag for epigenetic transcription activation (By similarity). Association with the 2-oxoglutarate dehydrogenase complex, which provides succinyl-CoA, is required for histone succinylation (By similarity). In different complexes, functions either as an acetyltransferase (HAT) or as a succinyltransferase: in the SAGA and ATAC complexes, acts as a histone acetyltransferase (By similarity). Has significant histone acetyltransferase activity with core histones, but not with nucleosome core particles (By similarity). Acetylation of histones gives a specific tag for epigenetic transcription activation (PubMed:28424240). Involved in long-term memory consolidation and synaptic plasticity: acts by promoting expression of a hippocampal gene expression network linked to neuroactive receptor signaling (PubMed:25024434). Acts as a positive regulator of T-cell activation: upon TCR stimulation, recruited to the IL2 promoter following interaction with NFATC2 and catalyzes acetylation of histone H3 at Lys-9 (H3K9ac), leading to promote IL2 expression (PubMed:28424240). Also acetylates non-histone proteins, such as CEBPB, PLK4 and TBX5 (PubMed:17301242). Involved in heart and limb development by mediating acetylation of TBX5, acetylation regulating nucleocytoplasmic shuttling of TBX5 (By similarity). Acts as a negative regulator of centrosome amplification by mediating acetylation of PLK4 (By similarity).[UniProtKB/Swiss-Prot Function]