

Product datasheet for MR222304L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Foxk1 (NM_199068) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Foxk1 (NM 199068) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Foxk

Synonyms: A630048H08Rik; Al463295; ENSMUSG00000075577; Gm10868; Mnf

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM_199068

ORF Size: 2157 bp

ORF Nucleotide

OTI Disclaimer:

_. __.

Sequence:

The ORF insert of this clone is exactly the same as(MR222304).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through paturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 199068.2, NP 951031.2

 RefSeq Size:
 7462 bp

 RefSeq ORF:
 2160 bp

 Locus ID:
 17425

 UniProt ID:
 P42128

Cytogenetics: 5 81.53 cM



Gene Summary:

Transcriptional regulator involved in different processes such as glucose metabolism, aerobic glycolysis, muscle cell differentiation and autophagy (PubMed:25402684, PubMed:29861159, PubMed:30700909). Recognizes and binds the forkhead DNA sequence motif (5'-GTAAACA-3') and can both act as a transcription activator or repressor, depending on the context (PubMed:25402684, PubMed:29861159, PubMed:30700909). Together with FOXK2, acts as a key regulator of metabolic reprogramming towards aerobic glycolysis, a process in which glucose is converted to lactate in the presence of oxygen (PubMed:30700909). Acts by promoting expression of enzymes for glycolysis (such as hexokinase-2 (HK2), phosphofructokinase, pyruvate kinase (PKLR) and lactate dehydrogenase), while suppressing further oxidation of pyruvate in the mitochondria by up-regulating pyruvate dehydrogenase kinases PDK1 and PDK4 (PubMed:30700909). Probably plays a role in gluconeogenesis during overnight fasting, when lactate from white adipose tissue and muscle is the main substrate (PubMed:30700909). Involved in mTORC1-mediated metabolic reprogramming: in response to mTORC1 signaling, translocates into the nucleus and regulates the expression of genes associated with glycolysis and downstream anabolic pathways, such as HIF1A, thereby regulating glucose metabolism (PubMed:29861159). Together with FOXK2, acts as a negative regulator of autophagy in skeletal muscle: in response to starvation, enters the nucleus, binds the promoters of autophagy genes and represses their expression, preventing proteolysis of skeletal muscle proteins (PubMed:25402684). Acts as a transcriptional regulator of the myogenic progenitor cell population in skeletal muscle (PubMed:8007964, PubMed:9271401, PubMed:12446708, PubMed:22956541). Binds to the upstream enhancer region (CCAC box) of myoglobin (MB) gene, regulating the myogenic progenitor cell population (PubMed:8007964, PubMed:9271401). Promotes muscle progenitor cell proliferation by repressing the transcriptional activity of FOXO4, thereby inhibiting myogenic differentiation (PubMed:12446708, PubMed:22956541). Involved in remodeling processes of adult muscles that occur in response to physiological stimuli (PubMed:9271401, PubMed:22956541). Required to correct temporal orchestration of molecular and cellular events necessary for muscle repair (PubMed:10792059). Represses myogenic differentiation by inhibiting MEFC activity (PubMed:22956541). Positively regulates Wnt/beta-catenin signaling by translocating DVL into the nucleus (By similarity). Reduces virus replication, probably by binding the interferon stimulated response element (ISRE) to promote antiviral gene expression (By similarity).[UniProtKB/Swiss-Prot Function]