

Product datasheet for MR222481L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Med1 (NM_001080118) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Med1 (NM_001080118) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Med1

Synonyms: Al480703; CRSP210; DRIP205; l11Jus15; PBP; Pparbp; TRAP220

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001080118

ORF Size: 4725 bp

ORF Nucleotide

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(MR222481).

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001080118.1</u>, <u>NP 001073587.1</u>

 RefSeq Size:
 6518 bp

 RefSeq ORF:
 4728 bp

 Locus ID:
 19014

 UniProt ID:
 Q925|9

Cytogenetics: 11 61.75 cM





Gene Summary:

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. Essential for embryogenesis, including development of the central nervous system, heart, liver and placenta and for erythropoiesis. Also required for normal transcriptional control of thyroid-stimulating hormone beta (TSHB) in the pituitary. Acts as a coactivator for GATA1-mediated transcriptional activation during erythroid differentiation of K562 erythroleukemia cells (By similarity).[UniProtKB/Swiss-Prot Function]