


Product datasheet for MR222633L3

Card9 (NM_001037747) Mouse Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids Product Name: Card9 (NM 001037747) Mouse Tagged Lenti ORF Clone Tag: Myc-DDK Symbol: Card9 Gm782 Synonyms: **Mammalian Cell** Puromycin Selection: Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092) E. coli Selection: Chloramphenicol (34 ug/mL) The ORF insert of this clone is exactly the same as(MR222633). **ORF** Nucleotide Sequence: **Restriction Sites:** Sgfl-Rsrll **Cloning Scheme:** Cloning sites used for ORF Shuttling: ORF Sqf I Rsr II ---- GCG ATC GC C ATG --- // --- NNN AG C GGA CCG --



ACCN: NM_001037747 ORF Size: 1608 bp

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	ard9 (NM_001037747) Mouse Tagged Lenti ORF Clone – MR222633L3
OTI Disclaimer:	Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <u>custsupport@origene.com</u> or by calling 301.340.3188 option 3 for pricing and delivery.
	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Me	 thod: 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 001037747.1</u>
RefSeq Size:	1611 bp
RefSeq ORF:	1611 bp
Locus ID:	332579
UniProt ID:	A2AIV8
Cytogenetics:	2 A3


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Card9 (NM_001037747) Mouse Tagged Lenti ORF Clone – MR222633L3

Gene Summary:

Adapter protein that plays a key role in innate immune response to a number of intracellular pathogens, such as C.albicans and L.monocytogenes. Is at the crossroads of ITAM-tyrosine kinase and the Toll-like receptors (TLR) and NOD2 signaling pathways (PubMed:17514206). Probably controls various innate immune response pathways depending on the intracellular pathogen. Controls CLEC7A (dectin-1)-mediated myeloid cell activation induced by the yeast cell wall component zymosan, leading to cytokine production and innate anti-fungal immunity: acts by regulating BCL10-MALT1-mediated NF-kappa-B activation pathway. Activates NF-kappa-B via BCL10 (PubMed:16862125). In response to the hyphal form of C.albicans, mediates CLEC6A (dectin-2)-induced I-kappa-B kinase ubiquitination, leading to NFkappa-B activation via interaction with BCL10 (PubMed:20538615). In response to L.monocytogenes infection, acts by connecting NOD2 recognition of peptidoglycan to downstream activation of MAP kinases (MAPK) without activating NF-kappa-B (PubMed:17187069). In response to fungal infection, may be required for the development and subsequent differentiation of interleukin 17-producing T helper (TH-17) cells (PubMed:17450144). Also involved in activation of myeloid cells via classical ITAM-associated receptors and TLR: required for TLR-mediated activation of MAPK, while it is not required for TLR-induced activation of NF-kappa-B (PubMed:17486093).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR222633L3

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US