

# Product datasheet for MR222682L4V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Mpp5 (NM\_019579) Mouse Tagged ORF Clone Lentiviral Particle

### **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** Mpp5 (NM\_019579) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Mpp5

**Synonyms:** 3830420B02Rik; Al255216; Al644496; Pals1

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_019579 **ORF Size:** 2028 bp

**ORF Nucleotide** 

2020 00

Sequence:

The ORF insert of this clone is exactly the same as(MR222682).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 019579.3, NP 062525.1

RefSeq Size: 5529 bp
RefSeq ORF: 2028 bp
Locus ID: 56217
UniProt ID: Q9|LB2
Cytogenetics: 12 C3







### **Gene Summary:**

Plays a role in tight junction biogenesis and in the establishment of cell polarity in epithelial cells (By similarity). Also involved in adherens junction biogenesis by ensuring correct localization of the exocyst complex protein EXOC4/SEC8 which allows trafficking of adherens junction structural component CDH1 to the cell surface (PubMed:17182851, PubMed:20237282). Plays a role through its interaction with CDH5 in vascular lumen formation and endothelial membrane polarity (By similarity). Required during embryonic and postnatal retinal development (PubMed:22398208). Required for the maintenance of cerebellar progenitor cells in an undifferentiated proliferative state, preventing premature differentiation, and is required for cerebellar histogenesis, fissure formation and cerebellar layer organization (PubMed:26657772). Plays a role in the radial and longitudinal extension of the myelin sheath in Schwann cells (PubMed:20237282). May modulate SC6A1/GAT1mediated GABA uptake by stabilizing the transporter (PubMed:15234345). May play a role in the T-cell receptor-mediated activation of NF-kappa-B (By similarity). Required for localization of EZR to the apical membrane of parietal cells and may play a role in the dynamic remodeling of the apical cytoskeleton (PubMed:15677456). Required for the normal polarized localization of the vesicular marker STX4 (PubMed:20237282). Required for the correct trafficking of the myelin proteins PMP22 and MAG (By similarity).[UniProtKB/Swiss-Prot Function]