

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR222714L3V

Ehmt2 (NM_145830) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Ehmt2 (NM_145830) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Ehmt2
Synonyms:	Bat8; D17Ertd710e; G9a; KMT1C; NG36
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_145830
ORF Size:	3792 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR222714).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 145830.1</u>
RefSeq Size:	4070 bp
RefSeq ORF:	3792 bp
Locus ID:	110147
UniProt ID:	<u>Q9Z148</u>
Cytogenetics:	17 18.45 cM



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Histone methyltransferase that specifically mono- and dimethylates 'Lys-9' of histone H3
(H3K9me1 and H3K9me2, respectively) in euchromatin. H3K9me represents a specific tag for
epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones. Also
mediates monomethylation of 'Lys-56' of histone H3 (H3K56me1) in G1 phase, leading to
promote interaction between histone H3 and PCNA and regulating DNA replication. Also
weakly methylates 'Lys-27' of histone H3 (H3K27me). Also required for DNA methylation, the
histone methyltransferase activity is not required for DNA methylation, suggesting that these
2 activities function independently. Probably targeted to histone H3 by different DNA-binding
proteins like E2F6, MGA, MAX and/or DP1. May also methylate histone H1. In addition to the
histone methyltransferase activity, also methylates non-histone proteins: mediates
dimethylation of 'Lys-373' of p53/TP53. Also methylates CDYL, WIZ, ACIN1, DNMT1, HDAC1,
ERCC6, KLF12 and itself.[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US