

Product datasheet for MR222733L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Copa (NM 009938) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Copa (NM 009938) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Copa

AU040324: xenin Synonyms:

Mammalian Cell

Selection:

ACCN:

None

NM 009938

Vector: pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag:

ORF Size: 3672 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR222733).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 009938.4, NP 034068.3

RefSeq Size: 4629 bp RefSeq ORF: 3675 bp Locus ID: 12847 **UniProt ID:** Q8CIE6

Cytogenetics: 1 79.54 cM







Gene Summary:

The coatomer is a cytosolic protein complex that binds to dilysine motifs and reversibly associates with Golgi non-clathrin-coated vesicles, which further mediate biosynthetic protein transport from the ER, via the Golgi up to the trans Golgi network. Coatomer complex is required for budding from Golgi membranes, and is essential for the retrograde Golgi-to-ER transport of dilysine-tagged proteins. In mammals, the coatomer can only be recruited by membranes associated to ADP-ribosylation factors (ARFs), which are small GTP-binding proteins; the complex also influences the Golgi structural integrity, as well as the processing, activity, and endocytic recycling of LDL receptors (By similarity).[UniProtKB/Swiss-Prot Function]