

Product datasheet for **MR222928**

Rpa1 (NM_001164223) Mouse Tagged ORF Clone

Product data:

| | |
|---------------------------|---|
| Product Type: | Expression Plasmids |
| Product Name: | Rpa1 (NM_001164223) Mouse Tagged ORF Clone |
| Tag: | Myc-DDK |
| Symbol: | Rpa1 |
| Synonyms: | 70kDa; 5031405K23Rik; AA589576; AW557552; RF-A; RP-A; Rpa |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-Entry (PS100001) |
| E. coli Selection: | Kanamycin (25 ug/mL) |



[View online »](#)

ORF Nucleotide
Sequence:

>MR222928 representing NM_001164223
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGGTGGGACACCTGAGCGAGGGGGCCATCGAGGTCATGATACAGCAGGAAAATACATCCATAAAGCCCA
TCCTGCAGGTCATTAACATCCGTCCCATTCTACAGGGAATAGGTCACCCCGGTACCGACTGCTCATGAG
TGATGGGTTGAACACGCTTTCCTCGTTCATGCTGGCCACGCAGCTGAACACCCTGGTCGAGGGAGGGCAG
CTGGCCAGCAACTGTGTCTGCCAGGTCACAAGTTCATCGTCAACACCCTGAAAGACGGCAGAATGCAAC
CAAAACCTCATGGGAATCCGTATGGTGAACAGTCAGTGTGAGGAAGTTAAGCCAGAAAGTAGTTGTTTT
GATGGACTTGAAGTTATGAAATCTGCTGAAGACGTTGGATTAAGATTGGGAATCCAGTGCCCTATAAT
GAAGGATATGGGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAAGCAGTCCCTTCCCAGCCTCTG
CAGCGACACCACCAGCCAGCAAGCCCCAGCCACAGAATGGGAGTCTGGGAATGGTTCCACTGCAGCTAA
GGCTTATGGTGCCTCAAAGCCATTTGAAAACCTGCCGGAAGTGGCTGTGCAGCCTTCAGGTGGGACA
CAGTCCAAAGTGGTGCCTCGCCATCGCCAGCCTCACCCCGTACCAAGTGGACTATTTGTCCCCGTGTCA
CCAACAAAAGTCAGATCCGCACCTGGAGTAATTCGCGGGGGAAGGAAAGCTTTCTCTCTAGAAGTGT
GGATGAAAGCGGTGAAATCAGAGCTACTGCTTTCAATGAGCAAGTGGACAAGTCTTTCCCTTATTGAA
GTGAACAAGGTGTATTACTTCTCAAAGGGCGCCCTGAAGATCGCTAACAAACAGTTCTCCGCTGTTAAAA
ATGACTATGAGATGACCTTCAATAATGAGACTTCTGTCTTCCCTGTGAAGATGGCCATCACTTACCCAC
AGTTCAGTTTGATTTACAGGGATCGGTGACCTAGAGAGCAAGGCTAAAGACGCACTAGTAGACATCATT
GGAATCTGCAAGAGCTATGAAGATTCGATTAATAACACAGTGAAGTGAACAACAGAGAAGTTGCTAAGA
GAAATATCTATTTGATGGACATGTCAGGGAAGGTGGTGACCACAACCTCTGTGGGAGAAGATGCTGACAA
GTTTGACGGCTCTCGGCAGCCCGTGTGGCCATCAAAGGAGCCGAGTCTCTGACTTCGGTGGACGGAGC
CTCTCGGTCTCTCATCCAGCACTGTCATTGTGAATCCTGACATCCCAGAGGCCTATAAGCTTCGTGGCT
GGTTTGACTCAGAAGGACAAGCCTTAGATGGTGTTCATCTCTGACCACAGGAGTGGAGGGGCCGGAGG
GGGCAACACCAACTGGAAAACCTTTCATGAGGCTAAATCTGAGAACCTGGCCAGGGAGACAAGGCGGAC
TATTTACGACTGTGGCAGCGGTGGTGTTCCTCGCAAAGAGAAGTGTGTACCAGGCCTGCCCAACCC
AGGACTGCAATAAGAAAGTATTGACCAGCAGAATGGGTGTACCCTGTGAGAAGTGTACCGGGAATT
TCCCAATTTCAAATACCGCATGATCTTATCGGCAATATTGCAGATTTTCAAGAGAACCAGTGGGTGACG
TGTTTCCAGGAGTCTGCAGAGGCCATCCTTGACAGAACTATGTACCTTGGGGAAGTCAAAGAGAAGA
ACGAGCAGGCGTTTGAAGAGTTTTCCAGAATGCCAAGTCCCGTCAATACGTTTCAAGATCAGGGTCAA
GCTGGAGACCTACAATGATGAATCTCGAATTAAGGCCACTGTGATGGACGTGAAGCCCGTGGACTTCAGA
GACTATGGCAGACGGCTGATCGGCAACATCAGGAAGAACATG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR222928 representing NM_001164223
 Red=Cloning site Green=Tags(s)

MVGHLSEGAIEVMIQQENTSIKPIIQVINIRPISTGNRSPRYRLLMSDGLNLTSSFMLATQLNLTVEGGQ
 LASNCVCQVHKFIVNTLKDGRMQPKPHGNPYGEQSVLRKLSQKVVVLMDLEVMSAEDVGLKIGNPVPYN
 EGYGQQQQQQQQQQVAVPSPASAATPPASKPQPQNGSLGMGSTAAKAYGASKPFGKPAAGTGLLQPSGGT
 QSKVVPIASLTPYQSKWTICARVTNKSQIRTWSNSRGEGLFSLLELVDESGEIRATAFNEQVDFKFFPLIE
 VNKVYYFSKGALKIANKQFSAVKNDYEMTFNNETSVLPCEDGHHLPTVQFDFGTGIGDLESKAKDALVDII
 GICKSYEDSIKITYKSNREVAKRNIYLMMSGKVVTTLWGEDADKFDGSRQPVMAIKGARVSDFGGRS
 LSVLSSSTVIIVNPDPEAYKLRGWFDFSEGQALDGVISIDHRSGGAGGGNTNWKTLHEAKSENLGQGDKAD
 YFSTVAVVFLRKENCMYQACPTQDCNKKVIDQQNGLYRCEKCDREFPNFKYRMLSANIADFQENQWWT
 CFQESAEAILGQNTMYLGELKEKNEQAFEEVFQANFRSFTFRIRVKLETYNDESRIKATVMDVKPVDFR
 DYGRRLIANIRKNM



TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/ja2721_g05.zip

Restriction Sites: Sgfl-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_001164223

ORF Size: 1935 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001164223.1](#), [NP_001157695.1](#)

RefSeq Size: 3121 bp

RefSeq ORF: 1935 bp

Locus ID: 68275


UniProt ID: [Q8VEE4](#)

Cytogenetics: 11 45.79 cM

MW: 71.4 kDa

Gene Summary:

As part of the heterotrimeric replication protein A complex (RPA/RP-A), binds and stabilizes single-stranded DNA intermediates, that form during DNA replication or upon DNA stress. It prevents their reannealing and in parallel, recruits and activates different proteins and complexes involved in DNA metabolism. Thereby, it plays an essential role both in DNA replication and the cellular response to DNA damage. In the cellular response to DNA damage, the RPA complex controls DNA repair and DNA damage checkpoint activation. Through recruitment of ATRIP activates the ATR kinase a master regulator of the DNA damage response. It is required for the recruitment of the DNA double-strand break repair factors RAD51 and RAD52 to chromatin in response to DNA damage. Also recruits to sites of DNA damage proteins like XPA and XPG that are involved in nucleotide excision repair and is required for this mechanism of DNA repair. Plays also a role in base excision repair (BER) probably through interaction with UNG. Also recruits SMARCAL1/HARP, which is involved in replication fork restart, to sites of DNA damage. May also play a role in telomere maintenance.[UniProtKB/Swiss-Prot Function]

Product images:


Circular map for MR222928