

Product datasheet for **MR223070L3V**

Tdg (NM_172552) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Tdg (NM_172552) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Tdg
Synonyms:	E130317C12Rik; JZA-3; Jza1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_172552
ORF Size:	1263 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR223070).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_172552.3
RefSeq Size:	3218 bp
RefSeq ORF:	1266 bp
Locus ID:	21665
UniProt ID:	P56581
Cytogenetics:	10 39.72 cM



[View online »](#)

Gene Summary:

DNA glycosylase that plays a key role in active DNA demethylation: specifically recognizes and binds 5-formylcytosine (5fC) and 5-carboxylcytosine (5caC) in the context of CpG sites and mediates their excision through base-excision repair (BER) to install an unmethylated cytosine (PubMed:21817016). Cannot remove 5-hydroxymethylcytosine (5hmC). According to an alternative model, involved in DNA demethylation by mediating DNA glycolase activity toward 5-hydroxymethyluracil (5hmU) produced by deamination of 5hmC (PubMed:21722948). Also involved in DNA repair by acting as a thymine-DNA glycosylase that mediates correction of G/T mismatches to G/C pairs: in the DNA of higher eukaryotes, hydrolytic deamination of 5-methylcytosine to thymine leads to the formation of G/T mismatches. Its role in the repair of canonical base damage is however minor compared to its role in DNA demethylation. It is capable of hydrolyzing the carbon-nitrogen bond between the sugar-phosphate backbone of the DNA and a mispaired thymine. In addition to the G/T, it can remove thymine also from C/T and T/T mismatches in the order G/T >> C/T > T/T. It has no detectable activity on apyrimidinic sites and does not catalyze the removal of thymine from A/T pairs or from single-stranded DNA. It can also remove uracil and 5-bromouracil from mismatches with guanine.[UniProtKB/Swiss-Prot Function]