

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR223071L3V

Pou4f2 (NM_138944) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Pou4f2 (NM_138944) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Pou4f2
Synonyms:	Brn-3.2; Brn-3b; Brn3b; mBrn3-3R; Pou4f-rs1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_138944
ORF Size:	1233 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR223071).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 138944.2</u>
RefSeq Size:	3212 bp
RefSeq ORF:	1236 bp
Locus ID:	18997
UniProt ID:	<u>Q63934</u>
Cytogenetics:	8 C1



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Gene Summary:

Tissue-specific DNA-binding transcription factor involved in the development and differentiation of target cells (PubMed:7904822, PubMed:8995448, PubMed:8972215, PubMed:10357904, PubMed:10414983, PubMed:11163266, PubMed:17668438, PubMed:25775587). Functions either as activator or repressor by modulating the rate of target gene transcription through RNA polymerase II enzyme in a promoter-dependent manner (PubMed:7904822, PubMed:7935408, PubMed:8065921, PubMed:7852360, PubMed:7797498, PubMed:8662774, PubMed:9694219, PubMed:10526314, PubMed:15733064, PubMed:17145718, PubMed:18368538). Binds to the consensus octamer motif 5'-AT[A/T]A[T/A]T[A/T]A-3' of promoter of target genes (PubMed:7904822, PubMed:8290353, PubMed:9111308, PubMed:10414983, PubMed:16152597, PubMed:17668438, PubMed:24643061). Plays a fundamental role in the gene regulatory network essential for retinal ganglion cell (RGC) differentiation (PubMed:8632990, PubMed:10357904, PubMed:25775587). Binds to an octamer site to form a ternary complex with ISL1; cooperates positively with ISL1 and ISL2 to potentiate transcriptional activation of RGC target genes being involved in RGC fate commitment in the developing retina and RGC axon formation and pathfinding (PubMed:8995448, PubMed:9261145, PubMed:8972215, PubMed:10357904, PubMed:11163266, PubMed:24643061, PubMed:25775587). Inhibits DLX1 and DLX2 transcriptional activities preventing DLX1- and DLX2-mediated ability to promote amacrine cell fate specification (PubMed:21875655). In cooperation with TP53 potentiates transcriptional activation of BAX promoter activity increasing neuronal cell apoptosis (PubMed:17145718). Negatively regulates BAX promoter activity in the absence of TP53 (PubMed:17145718). Acts as a transcriptional coactivator via its interaction with the transcription factor ESR1 by enhancing its effect on estrogen response element (ERE)containing promoter (PubMed:9448000). Antagonizes the transcriptional stimulatory activity of POU4F1 by preventing its binding to an octamer motif (PubMed:7935408, PubMed:8065921, PubMed:8537352, PubMed:7852360, PubMed:8662774). Involved in TNFSF11-mediated terminal osteoclast differentiation (PubMed:17668438).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US