

Product datasheet for MR223185L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cpeb3 (NM 198300) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Cpeb3 (NM_198300) Mouse Tagged ORF Clone Lentiviral Particle

Symbol:

4831444O18Rik; CPE-BP3; mKIAA0940 Synonyms:

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

mGFP Tag:

NM 198300 ACCN: **ORF Size:** 2148 bp

ORF Nucleotide

Sequence: OTI Disclaimer: The ORF insert of this clone is exactly the same as(MR223185).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 198300.3, NP 938042.2

RefSeq Size: 5792 bp RefSeq ORF: 2151 bp Locus ID: 208922 **UniProt ID:** Q7TN99 Cytogenetics: 19 C2







Gene Summary:

Sequence-specific RNA-binding protein which acts as a translational repressor in the basal unstimulated state but, following neuronal stimulation, acts as a translational activator (PubMed:17024188, PubMed:26074072). In contrast to CPEB1, does not bind to the cytoplasmic polyadenylation element (CPE), a uridine-rich sequence element within the mRNA 3' UTR, but binds to a U-rich loop within a stem-loop structure (PubMed:17024188). Required for the consolidation and maintenance of hippocampal-based long term memory (PubMed:26074003). In the basal state, binds to the mRNA 3' UTR of the glutamate receptors GRIA1 and GRIA2 and negatively regulates their translation (PubMed:17024188, PubMed:22153079). Also represses the translation of DLG4, GRIN1 GRIN2A and GRIN2B (PubMed:24155305). When activated, acts as a translational activator of GRIA1 and GRIA2 (PubMed:22153079, PubMed:26074003). In the basal state, suppresses SUMO2 translation but activates it following neuronal stimulation (PubMed:26074071). Binds to the 3' UTR of TRPV1 mRNA and represses TRPV1 translation which is required to maintain normal thermoception (PubMed:26915043). Binds actin mRNA, leading to actin translational repression in the basal state and to translational activation following neuronal stimulation (PubMed:26074072). Negatively regulates target mRNA levels by binding to TOB1 which recruits CNOT7/CAF1 to a ternary complex and this leads to target mRNA deadenylation and decay (By similarity). In addition to its role in translation, binds to and inhibits the transcriptional activation activity of STAT5B without affecting its dimerization or DNA-binding activity. This, in turn, represses transcription of the STAT5B target gene EGFR which has been shown to play a role in enhancing learning and memory performance (By similarity). In contrast to CPEB1, CPEB2 and CPEB4, not required for cell cycle progression (By similarity). [UniProtKB/Swiss-Prot Function]