

## Product datasheet for MR223387

### Cd247 (NM\_001113392) Mouse Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Cd247 (NM\_001113392) Mouse Tagged ORF Clone  
**Tag:** Myc-DDK  
**Symbol:** Cd247  
**Synonyms:** 4930549J05Rik; A430104F18Rik; AW552088; Cd3; Cd3-eta; Cd3-zeta; Cd3h; Cd3z; Cd3zeta; T3z; Tcrk  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**Cell Selection:** Neomycin  
**ORF Nucleotide Sequence:** >MR223387 representing NM\_001113392  
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGAAGTGAAAGTGTCTGTTCTCGCTGCATCCTCCACGTGCGGTTCCAGGAGCAGAGGCACAGAGCT  
TTGGTCTGCTGGATCCCAAACCTGCTACTTGTAGATGGAATCCTTTCATCTACGGAGTCATCATCAC  
AGCCCTGTACCTGAGAGCAAAATTCAGCAGGAGTGCAGAGACTGCTGCCAACCTGCAGGACCCCAACCAG  
CTCTACAATGAGCTCAATCTAGGGCGAAGAGAGGAATATGACGTCTTGGAGAAGAAGCGGGCTCGGGATC  
CAGAGATGGGAGGCAAACAGCAGAGGAGGAGGAACCCCGGAAGGCGTATACAATGCACCTGCAGAAAGA  
CAAGATGGCAGAAGCCTACAGTGAGATCGGCACAAAAGGCGAGAGGCGGAGGCAAGGGGCACGATGGC  
CTTTACCAGCCTGCCGACATAAGAAGCCTCTTAGCCTCCAGCAGCCGTATCCTAACACACTGTGCAGCA  
AAAACAGCTCTACAGGAGAGGAAGGAGATGTCATTGCGCTG

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >MR223387 representing NM\_001113392  
Red=Cloning site Green=Tags(s)


MKWKVSVLACILHVRFPGAEAQSFGLLDPKLCYLLDGILFIYGVIIITALYLRAKFSRSAETAANLQDPNQ  
LYNELNLGRREYDVLEKKRRARDPEMGGKQRRRNPQEGVYNALQKDKMAEAYSEIGTKGERRRGKGHG  
LYQPADIRSLASQPPYNTLCSKNSSTGEEGDVIRL

**TR**TRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:** Sgfl-MluI



**Cloning Scheme:**

**Plasmid Map:**


**ACCN:** NM\_001113392

**ORF Size:** 531 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u>NM_001113392.2, NP_001106863.1</u>
<b>RefSeq Size:</b>	5538 bp
<b>RefSeq ORF:</b>	534 bp
<b>Locus ID:</b>	12503
<b>Cytogenetics:</b>	1 73.14 cM
<b>MW:</b>	20.5 kDa
<b>Gene Summary:</b>	<p>Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways. CD3Z ITAMs phosphorylation creates multiple docking sites for the protein kinase ZAP70 leading to ZAP70 phosphorylation and its conversion into a catalytically active enzyme. Plays an important role in intrathymic T-cell differentiation. Additionally, participates in the activity-dependent synapse formation of retinal ganglion cells (RGCs) in both the retina and dorsal lateral geniculate nucleus (dLGN). [UniProtKB/Swiss-Prot Function]</p>