

Product datasheet for MR223614L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Slc8b1 (NM_001177595) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Slc8b1 (NM_001177595) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Slc8b1

Synonyms: AF261233; NCKX6; NCLX; Slc24a6

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001177595

ORF Size: 1656 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR223614).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001177595.1, NP 001171066.1

RefSeq Size: 2725 bp
RefSeq ORF: 1659 bp
Locus ID: 170756
UniProt ID: Q925Q3

Cytogenetics: 5 F





Gene Summary:

Mitochondrial sodium/calcium antiporter that mediates sodium-dependent calcium efflux from mitochondrion, by mediating the exchange of 3 sodium ions per 1 calcium ion (PubMed:20018762, PubMed:28445457). Plays a central role in mitochondrial calcium homeostasis by mediating mitochondrial calcium extrusion: calcium efflux is essential for mitochondrial function and cell survival, notably in cardiomyocytes (PubMed:24067497, PubMed:28445457). Regulates rates of glucose-dependent insulin secretion in pancreatic beta-cells during the first phase of insulin secretion: acts by mediating efflux of calcium from mitochondrion, thereby affecting cytoplasmic calcium responses (By similarity). Required for store-operated Ca(2+) entry (SOCE) and Ca(2+) release-activated Ca(2+) (CRAC) channel regulation: sodium transport by SLC8B1 leads to promote calcium-shuttling that modulates mitochondrial redox status, thereby regulating SOCE activity (By similarity). Involved in B-lymphocyte chemotaxis (PubMed:27328625). Able to transport Ca(2+) in exchange of either Li(+) or Na(+), explaining how Li(+) catalyzes Ca(2+) exchange (By similarity). In contrast to other members of the family its function is independent of K(+) (By similarity). [UniProtKB/Swiss-Prot Function]