

## **Product datasheet for MR223685L3**

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

## Brsk2 (NM\_001009930) Mouse Tagged Lenti ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

Product Name: Brsk2 (NM\_001009930) Mouse Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: Brsk2

Synonyms: 4833424K13Rik; SAD-A; SADA

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)


ORF Nucleotide The ORF insert of this clone is exactly the same as(MR223685).

Sequence:

Restriction Sites: Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF.

**ACCN:** NM\_001009930

ORF Size: 2157 bp



### Brsk2 (NM\_001009930) Mouse Tagged Lenti ORF Clone - MR223685L3

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001009930.3</u>, <u>NP 001009930.1</u>

 RefSeq Size:
 2160 bp

 RefSeq ORF:
 2160 bp

 Locus ID:
 75770

 UniProt ID:
 Q69Z98

 Cytogenetics:
 7 F5

**Gene Summary:** Serine/threonine-protein kinase that plays a key role in polarization of neurons and

axonogenesis, cell cycle progress and insulin secretion. Phosphorylates CDK16, CDC25C, MAPT/TAU, PAK1 and WEE1. Following phosphorylation and activation by STK11/LKB1, acts as a key regulator of polarization of cortical neurons, probably by mediating phosphorylation of microtubule-associated proteins such as MAPT/TAU at 'Thr-504' and 'Ser-554'. Also regulates neuron polarization by mediating phosphorylation of WEE1 at 'Ser-642' in post-mitotic neurons, leading to down-regulate WEE1 activity in polarized neurons. Plays a role in the regulation of the mitotic cell cycle progress and the onset of mitosis. Plays a role in the regulation of insulin secretion in response to elevated glucose levels, probably via phosphorylation of CDK16 and PAK1. While BRSK2 phosphorylated at Thr-175 can inhibit insulin secretion (PubMed:22798068), BRSK2 phosphorylated at Thr-261 can promote insulin


secretion (PubMed:22669945). Regulates reorganization of the actin cytoskeleton. May play a

role in the apoptotic response triggered by endoplasmic reticulum (ER) stress.

[UniProtKB/Swiss-Prot Function]



# **Product images:**



Circular map for MR223685L3