

Product datasheet for MR223690L3


Brd8 (NM_030147) Mouse Tagged Lenti ORF Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type: | Expression Plasmids |
|------------------------------|---|
| Product Name: | Brd8 (NM_030147) Mouse Tagged Lenti ORF Clone |
| Tag: | Myc-DDK |
| Symbol: | Brd8 |
| Synonyms: | 2610007E11Rik; 4432404P07Rik; SMAP |
| Mammalian Cell Selection: | Puromycin |
| Vector: | pLenti-C-Myc-DDK-P2A-Puro (PS100092) |
| E. coli Selection: | Chloramphenicol (34 ug/mL) |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(MR223690). |
| Restriction Sites: | Sgfl-Mlul |
| Cloning Scheme: | Cloning sites used for ORF Shuttling: |
| | Sgf1 ORF Mlu1 GCG ATC GC ATG// NNÑ ACG CGT |



ACCN: ORF Size: NM_030147 2853 bp



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


| DRIGENE Brd8 (NM_030147) Mouse Tagged Lenti ORF Clone – MR223690L3 | |
|---|--|
| OTI Disclaimer: | Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <u>custsupport@origene.com</u> or by calling 301.340.3188 option 3 for pricing and delivery. |
| | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u> |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| Components: | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water). |
| Reconstitution Met | 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | <u>NM 030147.2</u> |
| RefSeq Size: | 4677 bp |
| RefSeq ORF: | 2856 bp |
| Locus ID: | 78656 |
| UniProt ID: | <u>Q8R3B7</u> |
| Cytogenetics: | 18 B1 |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Brd8 (NM_030147) Mouse Tagged Lenti ORF Clone – MR223690L3

Gene Summary:May act as a coactivator during transcriptional activation by hormone-activated nuclear
receptors (NR). Stimulates transcriptional activation by AR/DHTR, ESR1/NR3A1, RXRA/NR2B1
and THRB/ERBA2. Component of the NuA4 histone acetyltransferase (HAT) complex which is
involved in transcriptional activation of select genes principally by acetylation of nucleosomal
histones H4 and H2A. This modification may both alter nucleosome - DNA interactions and
promote interaction of the modified histones with other proteins which positively regulate
transcription. This complex may be required for the activation of transcriptional programs
associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor
mediated growth arrest and replicative senescence, apoptosis, and DNA repair. NuA4 may
also play a direct role in DNA repair when recruited to sites of DNA damage. Component of a
SWR1-like complex that specifically mediates the removal of histone H2A.Z/H2AFZ from the
nucleosome.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR223690L3

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US