

Product datasheet for MR223868

Max (NM 008558) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Max (NM_008558) Mouse Tagged ORF Clone

Tag: Myc-DDK

Symbol: Max

Synonyms: AA960152; Al875693; bHLHd4; bHLHd5; bHLHd6; bHLHd7; bHLHd8

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>MR223868 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGAGCGATAACGATGACATCGAGGTGGAGAGCGACGAAGAGCAACCGAGGTTTCAATCTGCGGCTGACA
AGCGGGCTCACCATAATGCACTGGAACGAAAACGTAGGGACCACATCAAAGACAGCTTTCACAGTTTGCG
GGACTCAGTCCCATCACTCCAAGGAGAAGAGCATCCCGGGCCCAAATCCTAGACAAAGCAACAGAGTAT
ATCCAGTATATGCGAAGGAAAAACCATACGCACCAGCAAGACATTGATGACCTCAAGCGGCAGAATGCTC
TTCTGGAGCAACAAGTCCGTGCACTGGAGAAGGCAAGATCAAGTGCCCAACTGCAGACCAACTACCCCTC
CTCAGACCAACAGCCTCTACACCCAACGCCAAGGGCGGCACCATCTCTGCCTTCGATGGGGGTTCAGACTCC

AGCTCAGAATCCGAGCCTGAAGAGCCCCAGAGCAGGAAGAAACTCCGGATGGAGGCCAGC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAG**GTTTAA**

Protein Sequence: >MR223868 protein sequence

Red=Cloning site Green=Tags(s)

MSDNDDIEVESDEEQPRFQSAADKRAHHNALERKRRDHIKDSFHSLRDSVPSLQGEKASRAQILDKATEY IQYMRRKNHTHQQDIDDLKRQNALLEQQVRALEKARSSAQLQTNYPSSDNSLYTNAKGGTISAFDGGSDS

SSESEPEEPQSRKKLRMEAS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-Mlul




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_008558

ORF Size: 480 bp

OTI Disclaimer:

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at customport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

OTI Annotation:

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.



Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 008558.2, NP 032584.2</u>

 RefSeq Size:
 2005 bp

 RefSeq ORF:
 483 bp

 Locus ID:
 17187

 UniProt ID:
 P28574


Cytogenetics: 12 33.78 cM **MW:** 18.2 kDa

Gene Summary: Transcription regulator. Forms a sequence-specific DNA-binding protein complex with MYC or

MAD which recognizes the core sequence 5'-CAC[GA]TG-3'. The MYC:MAX complex is a transcriptional activator, whereas the MAD:MAX complex is a repressor. CpG methylation of the recognition site greatly inhibits DNA binding, suggesting that DNA methylation may regulate the MYC:MAX complex in vivo. May repress transcription via the recruitment of a chromatin remodeling complex containing H3 'Lys-9' histone methyltransferase activity. Represses MYC transcriptional activity from E-box elements (By similarity).[UniProtKB/Swiss-

Prot Function]

Product images:



Circular map for MR223868