

Product datasheet for MR224028L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Pou2f2 (NM_001163554) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Pou2f2 (NM_001163554) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Pou2f2

Synonyms: Oct-2; Oct2a; Oct2b; Oct2c; Oct2d; Otf-2; Otf2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001163554

ORF Size: 1749 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR224028).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001163554.1</u>, <u>NP 001157026.1</u>

 RefSeq Size:
 3323 bp

 RefSeq ORF:
 1752 bp

 Locus ID:
 18987

 UniProt ID:
 Q00196

 Cytogenetics:
 7 13.73 cM







Gene Summary:

Transcription factor that specifically binds to the octamer motif (5'-ATTTGCAT-3'). Regulates transcription in a number of tissues in addition to activating immunoglobulin gene expression. Modulates transcription transactivation by NR3C1, AR and PGR. Isoform OCT2.5 activates the U2 small nuclear RNA (snRNA) promoter. Isoforms OCT2.1, OCT2.2 and OCT2.3 activate octamer-containing promoters. Isoforms OCT2.4 and OCT2.5 repress some promoters and activate others. Isoform OCT2.7 is unable to bind to the octamer motif, but can still activate the beta-casein gene promoter at low levels.[UniProtKB/Swiss-Prot Function]