

## Product datasheet for MR224137L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Adamts12 (NM\_175501) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** Adamts12 (NM\_175501) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Adamts12

Synonyms: ADAMTS-12; Al605170

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_175501 **ORF Size:** 4800 bp

**ORF Nucleotide** 

\_\_\_\_\_

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(MR224137).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 175501.3, NP 780710.2

RefSeq Size: 8622 bp
RefSeq ORF: 4803 bp
Locus ID: 239337
UniProt ID: Q811B3
Cytogenetics: 15 A1







## **Gene Summary:**

This gene encodes a member of "a disintegrin and metalloproteinase with thrombospondin motifs" (ADAMTS) family of multi-domain matrix-associated metalloendopeptidases that have diverse roles in tissue morphogenesis and pathophysiological remodeling, in inflammation and in vascular biology. The encoded preproprotein undergoes proteolytic processing to generate an active protease. Mice lacking the encoded protein exhibit increased angiogenic response and tumor invasion in different models of angiogenesis and, severe inflammation and delayed recovery when subjected to experimental conditions that induce colitis, endotoxic sepsis and pancreatitis. [provided by RefSeq, Jul 2016]