

Product datasheet for **MR224240**

Golga2 (NM_001080968) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Golga2 (NM_001080968) Mouse Tagged ORF Clone
Tag: Myc-DDK
Symbol: Golga2
Synonyms: AW555139; GM130; mKIAA4150
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
ORF Nucleotide Sequence: >MR224240 representing NM_001080968
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**GCGATCGCC**

ATGTGGCCCCCGCTTCCCCCTCCCCGCCCGGGATGTCGGAAGAAACCAGGCAGAGCAAATGGCTG
CGGCCAAGAAAAAGCTGAGAGAGTATCAGCAGAAGAACAGCCCGGTGTTCTGCAGGAGCAAAGAAGAA
GAAAAAGATTAATAATGGCCATAGCCCAGAGAGACCCACTGCCAGTGACTGTCAGTCACCAGAGAATGTG
CCACAGACCATATTGCTCCTGCACCGCAACCGCTGCTACTGACACTATGTTTCTTGGTGTACCCCTT
CCCTGATGCTGACCTCACTCAGAGCCATGATGCTGGCAATTGCTCTAACCTCATGGAGGAGACCAAGAC
TTTCTCATCGACTGAGAGTCTGCGACAACCTTTCTCAACAGCTCAATGGCCTTGTGTCTGAGTCTACATCC
TACATCAACGGGGAGGGCCTCACATCTTCCAACATGAAGGAACTGGAGAACCAGGTACCAAGAGCTAGCAG
TAGCCCTGGACTCCAGCTATGTAACAAACAAACAACACTCAGTAGCACGATAGAGGAATTGAAACAACAGAA
CCAAGACACTCTGGATCAACTGGAGAAAGAGAAGAAGGATTATCAGCAGAAGCTGGCAAAGGAGCAGGGC
TCTCTCAGGGAACAGCTGCAGGTTACATTCAGACCATAGGCATCCTGGTGTCTGAGAAGGCAGAAATTAC
AGACGGCCCTGGCCACACTCAGCAAGCAGCCAGGCAGAAAGCAGGAGAGTCCGAGGATCTTGCCAGCCC
ACTACAGTCTCCCGACAGCGTGTGGGAGAGCTGGAGCGGACACTGTCTACCGTGTCAACGCAACAGAA
CAGGCAGACAGGTATAACAAGGACCTAACCAAAGAGCGGGATGCCCTCAAGCTGGAGCTGTATAAGAACA
GCAAAAGCAATGAGGACCTGAGGCAGCAGAACTCAGAACTGGAAGAGAACTCCGAGTCTGGTGGCAGA
GAAAGCAGCTGCACAGTTGGGGTGGAGGAGCTGCAGAAGAAGTTAGAGATGTCAGAGCTGCTGCTGCAG
CAGTTTTCTAGTCAGTCTTCGGCCGAGGCGGTAATGAGCAGTTACAACATGCCATGGAGGAGCGGGCTC
AGCTGGAGACCCATGTTAGCCAGCTGATGGAATCACTGAAGCAGCTCCAGGTGGAGAGAGACCAGTATGC
GGAGAACCTAAAAGGAGAGAGTGCCATGTGGCAGCAGAGGTACAGCAAATGGCCGAGCAGGTGCATACA
CTGAAGGAAGAGAAGGAGCACAGAGAACGCAGGTACAAGAGCTGGAGACCAGCTTGCCGCACTCCGGA
GTCAGATGGAGGAGCCACCGCTCCAGAGCCTCCAGCTGGGCCCTCCGAGCGGAAGAGCAGCTGCAGGG
AGAGGTCGAGCAGCTGCAAGGAACTGGAGAGGCTGACGGGACAGCTACGGGCTCAGGTGCAGGACAAT
GAGAGCCTGAGCCACCTCAACCGGGAGCAGGAGGGGCTGCTGCTGGAGCTGGAGCGGGAGGCCAGCGCT



GGAGCGAGCAGGCCGAGGAGCGCAAGCAGATCCTGGAGAGCATGCAGAGTGACCGCACCACCATCAGCAG
 AGCACTGTACAGAACCCGAGAGCTGAAGGAGCAGCTGGCCGAGCTGCAGAACGGCTTCGTGAGGCTGACC
 AATGAGAACATGGAGATCACCAGTGCAGTGCAGTGCAGAGCAGCATGTCAAGAAAGAGCTAGCCCGGAAGC
 TGGGAGAGCTGCAGGAGAGACTGGGAGAGCTGAAGGAGACGGTGGAGCTCAAGAGCCAAGAAGCACAGGG
 TCTACAGGAGCAGCGAGACCAGTGCCTGTCCCACCTGCAGCAGTACGCTGCTGCTACCAACAGCACTTG
 GCGGCCTATGAGCAGCTGACCTCTGAGAAGGAGGCAATACACAAGCAGCTTCTGCTGCAGACTCAGCTCA
 TGGACCAGCTGCAGCATGAGGAGGTACAGGGCAAGATGGCAGCCGAGCTGGCCCGCCAGGAGCTGCAGGA
 GGCCCGAGGAGCGCTAAAAGCTACCAGCCAGGAGAACCAACAGCTCCAAGCCAGCTGAGCCTCTTGCTC
 CTCCCTGGGAAGGAGATGTGGACCAGGAGGAAGAAGATGAGGAGGTTCCCTCAGTCGAGTCTGGCCATCC
 CAGAAGACCTAGACAGTAGAGAGGCCATGGTGGCATTCTTTAATGCCGCTATAGCCAGAGCAGAGGAAGA
 ACAGGCCCGACTACGTGTGCAGCTGAAAGAGCAGAAGGCACGTTGCCGGAGTTTGTCTCACTTGGCAGCC
 CCAGTCCAAGCAAGCTTAAAAGGAGGCAGTGGTCCCCAGGAATGTGGATGACTCTGCATCTGAGGAGA
 GCAATCAGGCCCTACATGTAGCCATGGAGAAGCTGCAGAGCCGCTTCTGGAGGTCATGCAGGAGAAAGT
 GGAGCTCAAGGAGAGGGTAGAGGAGCTTGAACATTGCTGCATTACGCTTCTGGAGAGACAGACACCATT
 GGAGAGTACATTGCCCTTACAAAATCAGAGGGCGGTGCTGAAGGCTCGGCATTTGGAGAAGGAAGAAT
 ACATCAGCCGGCTGGCTCAGGACAAGGAGGAGATGAAGGTGAAGCTGCTGGAGCTCCAGGAGCTGGTGT
 GAGGCTTGTGAACGAGCGCAATGAATGGCAGGGCAAGTTCCTGGCCGCTCTCAGAACCCTGGTGACGTG
 CTCACCCAGTGGCCACAGGCTCCCAGGAATTTGGTGTGCTGACCAGCAGGATGACCTCAGGGAGGTGA
 GCCTGGCTGACGATATAGAGCCTGCACAAGGAGAGGCAGGGGTACCTGCTCCCCATGAGAACCCTACTGC
 ACAGCAGATCATGCAGCTGCTGCGTGAGATCCAGAACCCCGGGAGCGGCCAGGCCTGGGTAGCAACCTT
 TGCATCCCCTTTTCTACCGTCCGATGAGAACGACGAGGTGAAGATCATGGTTGTA

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence:

>MR224240 representing NM_001080968
 Red=Cloning site Green=Tags(s)

MWPPRFPPPRPGMSEETRQSKLAAAKKKLREYQKNSPGVPAAGAKKKKIKNGHSPERPTASDCQSPENV
 PTDHIAPAPPTAATDTMFLGVTPSPDADLTQSHDAGNCSNLMEETKTFSSTESLRQLSQQLNGLVSESTS
 YINGEGLTSSNMKELENRYQELAVALDSSYVTNKQLSSTIEELKQQNQDLDQLEKEKKDYQQLAKEQG
 SLREQLQVHIQTIGILVSEKAEALQALHTQQAARQKAGESEDLASRLQSSRQRVGELERTLSTVSTQQK
 QADRYNKDLTKERDALKLELYKNSKSNEDLRQNSELEEKLRLVVAEKAAQLGVEELQKKLEMSELLLQ
 QFSSQSSAAGGNEQLQHAMEERAQLETHVSQLMESLKQLQVERDQYAENLKGESAMWQQRVQMAEQVHT
 LKEEKEHRERQVQLELTSLAALRSQMEEPPPEPPAGPSEAEELQGEVEQLHKELERLTGQLRAQVQDN
 ESLSHLNREQEGRLLLELERAQRWSEQAERKQILESMQSDRTTISRALSQNRELKEQLAELQNGFVRLT
 NENMEITSALQSEQHVKKELARKLGELQERLGELKVELKSQEAQGLQEQRDQCLSHLQQYAAAQQHL
 AAAYEQLTSEKEAIHKQLLLQTQLMDQLQHEEVQGMMAELARQELQEAQERLKATSQENQQLQAQLSLLV
 LPGEGDQVEEEDVEPQSSLAIPEDLDSREAMVAFNAAIARAEQARLRVQLKEQKARCRSLSHLAA
 PVQSKLEKEAVVPRNVDDASSEESNQALHVAMEKLSRFLEVMQEKVELKERVEELEHCCIQLSGETDTI
 GEYIALYQNRVLAHLEKKEEYISRLAQDKEEMKVKLLELQELVLRVNERNEWQKFLAVSQNPQGDV
 LTPVPTGSQEFGAADQDDLREVSLADDIEPAQGEAGVPAPHENPTAQQIMQLLREIQNPRERPLGNSP
 CIPFFYRADENDEVKIMVV


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

Sgfl-Mlul


Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

Plasmid Map:



ACCN: NM_001080968
 ORF Size: 2997 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001080968.1 , NP_001074437.1
RefSeq Size:	4396 bp
RefSeq ORF:	3000 bp
Locus ID:	99412
UniProt ID:	Q921M4
Cytogenetics:	2 B
MW:	113.7 kDa

Gene Summary:

Peripheral membrane component of the cis-Golgi stack that acts as a membrane skeleton that maintains the structure of the Golgi apparatus, and as a vesicle tether that facilitates vesicle fusion to the Golgi membrane (PubMed:28028212). Required for normal protein transport from the endoplasmic reticulum to the Golgi apparatus and the cell membrane (PubMed:28028212). Together with p115/USO1 and STX5, involved in vesicle tethering and fusion at the cis-Golgi membrane to maintain the stacked and inter-connected structure of the Golgi apparatus. Plays a central role in mitotic Golgi disassembly: phosphorylation at Ser-37 by CDK1 at the onset of mitosis inhibits the interaction with p115/USO1, preventing tethering of COPI vesicles and thereby inhibiting transport through the Golgi apparatus during mitosis. Also plays a key role in spindle pole assembly and centrosome organization (By similarity). Promotes the mitotic spindle pole assembly by activating the spindle assembly factor TPX2 to nucleate microtubules around the Golgi and capture them to couple mitotic membranes to the spindle: upon phosphorylation at the onset of mitosis, GOLGA2 interacts with importin-alpha via the nuclear localization signal region, leading to recruit importin-alpha to the Golgi membranes and liberate the spindle assembly factor TPX2 from importin-alpha. TPX2 then activates AURKA kinase and stimulates local microtubule nucleation. Upon filament assembly, nascent microtubules are further captured by GOLGA2, thus linking Golgi membranes to the spindle (By similarity). Regulates the meiotic spindle pole assembly, probably via the same mechanism (PubMed:21552007). Also regulates the centrosome organization (By similarity). Also required for the Golgi ribbon formation and glycosylation of membrane and secretory proteins (By similarity).[UniProtKB/Swiss-Prot Function]