


## Product datasheet for MR224582

### Foxe3 (NM\_015758) Mouse Tagged ORF Clone

#### Product data:

<b>Product Type:</b>	Expression Plasmids
<b>Tag:</b>	Myc-DDK
<b>Symbol:</b>	Foxe3
<b>Synonyms:</b>	dyl; FREAC8; rct
<b>Mammalian Cell Selection:</b>	Neomycin
<b>Vector:</b>	pCMV6-Entry (PS100001)
<b>E. coli Selection:</b>	Kanamycin (25 ug/mL)
<b>Restriction Sites:</b>	SgfI-MluI
<b>Cloning Scheme:</b>	

Cloning sites used for ORF Shuttling:




\* The last codon before the Stop codon of the ORF

<b>ACCN:</b>	NM_015758
<b>ORF Size:</b>	864 bp



<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>Note:</b>	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
<b>RefSeq:</b>	<a href="#">NM_015758.3</a>
<b>RefSeq Size:</b>	867 bp
<b>RefSeq ORF:</b>	867 bp
<b>Locus ID:</b>	30923
<b>UniProt ID:</b>	<a href="#">Q9QY14</a>
<b>Cytogenetics:</b>	4 D1
<b>MW:</b>	30.5 kDa
<b>Gene Summary:</b>	Transcription factor that controls lens epithelial cell growth through regulation of proliferation, apoptosis and cell cycle (PubMed:10652278, PubMed:10890982). During lens development, controls the ratio of the lens fiber cells to the cells of the anterior lens epithelium by regulating the rate of proliferation and differentiation (PubMed:16199865). Controls lens vesicle closure and subsequent separation of the lens vesicle from ectoderm (PubMed:10652278). Is required for morphogenesis and differentiation of the anterior segment of the eye (PubMed:17064680). Controls the expression of DNAJB1 in a pathway that is crucial for the development of the anterior segment of the eye (By similarity).[UniProtKB/Swiss-Prot Function]

## Product images:



Circular map for MR224582