

Product datasheet for MR224590L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Pcgf5 (NM_029508) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Pcgf5 (NM 029508) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Pcgf5

Synonyms: 0610009F02Rik; 1110054A01Rik; 5830406C17Rik; 5830443C21Rik; 9530023M17Rik; Al324127

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_029508

ORF Size: 708 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR224590).

OTI Disclaimer:

Cytogenetics:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 029508.3

 RefSeq Size:
 1509 bp

 RefSeq ORF:
 711 bp

 Locus ID:
 76073

 UniProt ID:
 Q3UK78

19 C2







Gene Summary:

Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility (PubMed:27136092, PubMed:28596365). Within the PRC1-like complex, regulates RNF2 ubiquitin ligase activity (By similarity). Plays a redundant role with PCGF3 as part of a PRC1-like complex that mediates monoubiquitination of histone H2A 'Lys-119' on the X chromosome and is required for normal silencing of one copy of the X chromosome in XX females (PubMed:28596365).[UniProtKB/Swiss-Prot Function]