

## Product datasheet for MR224833L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## P3h3 (NM\_013534) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** P3h3 (NM\_013534) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: P3h3

Synonyms: BC016431; Grcb; Leprel2

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_013534

ORF Size: 2196 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR224833).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 013534.4

 RefSeq Size:
 2779 bp

 RefSeq ORF:
 2199 bp

 Locus ID:
 14789

 UniProt ID:
 Q8CG70

 Cytogenetics:
 6 59.17 cM







## **Gene Summary:**

Part of a complex composed of PLOD1, P3H3 and P3H4 that catalyzes hydroxylation of lysine residues in collagen alpha chains and is required for normal assembly and cross-linkling of collagen fibrils (PubMed:27119146). Required for normal hydroxylation of lysine residues in type I collagen chains in skin, bone, tendon, aorta and cornea (PubMed:28115524). Required for normal skin stability via its role in hydroxylation of lysine residues in collagen alpha chains and in collagen fibril assembly (PubMed:27119146, PubMed:28115524). Apparently not required for normal prolyl 3-hydroxylation on collagen chains, possibly because it functions redundantly with other prolyl 3-hydroxylases (PubMed:28115524). [UniProtKB/Swiss-Prot Function]