

## Product datasheet for **MR224847L3V**

### Deaf1 (NM\_016874) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Deaf1 (NM_016874) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Deaf1
Synonyms:	AU042387; C230009B13Rik; NUDR
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_016874
ORF Size:	1698 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR224847).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_016874.2</a> , <a href="#">NP_058570.1</a>
RefSeq Size:	2192 bp
RefSeq ORF:	1701 bp
Locus ID:	54006
UniProt ID:	<a href="#">Q9Z1T5</a>
Cytogenetics:	7 F5



[View online »](#)

**Gene Summary:**

Transcription factor that binds to sequence with multiple copies of 5'-TTC[CG]G-3' present in its own promoter and that of the HNRPA2B1 gene. Down-regulates transcription of these genes. Binds to the retinoic acid response element (RARE) 5'-AGGGTTCACCGAAAGTTCA-3'. Activates the proenkephalin gene independently of promoter binding, probably through protein-protein interaction (By similarity). Regulates epithelial cell proliferation and side-branching in the mammary gland. Required for neural tube closure and skeletal patterning. Controls the expression of peripheral tissue antigens in pancreatic lymph nodes. Isoform 1 displays greater transcriptional activity than isoform 2. Isoform 2 may inhibit transcriptional activity of isoform 1 by interacting with it and retaining it in the cytoplasm. Transcriptional activator of EIF4G3 (By similarity). May also involved in behavior (PubMed:24726472). [UniProtKB/Swiss-Prot Function]