

Product datasheet for **MR224900L4V**

Asah2 (NM_018830) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Asah2 (NM_018830) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Asah2
Synonyms:	AI585898
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_018830
ORF Size:	2268 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR224900).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_018830.1 , NP_061300.1
RefSeq Size:	4835 bp
RefSeq ORF:	2271 bp
Locus ID:	54447
UniProt ID:	Q9JHE3
Cytogenetics:	19 C1



[View online »](#)

Gene Summary:

Plasma membrane ceramidase that hydrolyzes sphingolipid ceramides into sphingosine and free fatty acids at neutral pH (PubMed:10753931, PubMed:10652340, PubMed:16380386). Ceramides, sphingosine, and its phosphorylated form sphingosine-1-phosphate are bioactive lipids that mediate cellular signaling pathways regulating several biological processes including cell proliferation, apoptosis and differentiation (PubMed:14557071). Also catalyzes the reverse reaction allowing the synthesis of ceramides from fatty acids and sphingosine (PubMed:10652340, PubMed:21613224). Together with sphingomyelinase, participates in the production of sphingosine and sphingosine-1-phosphate from the degradation of sphingomyelin, a sphingolipid enriched in the plasma membrane of cells (PubMed:16126722). Also participates in the hydrolysis of ceramides from the extracellular milieu allowing the production of sphingosine-1-phosphate inside and outside cells (PubMed:16126722). This is the case for instance with the digestion of dietary sphingolipids in the intestinal tract (PubMed:16380386).[UniProtKB/Swiss-Prot Function]