

Product datasheet for **MR224932**

Cd247 (NM_001113394) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Cd247 (NM_001113394) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Cd247
Synonyms:	4930549J05Rik; A430104F18Rik; AW552088; Cd3; Cd3-eta; Cd3-zeta; Cd3h; Cd3z; Cd3zeta; T3z; Tcrk
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR224932 representing NM_001113394 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**GCGATCGCC**

ATGAAGTGGAAAGTGTCTGTTCTCGCCTGCATCCTCCACGTGCGGTTCCAGGAGCAGAGGCACAGAGCT
TTGGTCTGCTGGATCCCAAACCTGCTACTTGCTAGATGGAATCCTTTCATCTACGGAGTCATCATCAC
AGCCCTGTACCTGAGAGCAAAATTCAGCAGGAGTGCAGAGACTGCTGCCAACCTGCAGGACCCCAACCAG
CTCTACAATGAGCTCAATCTAGGGCGAAGAGAGGAATATGACGTCTTGGAGAAGAAGCGGGCTCGGGATC
CAGAGATGGGAGGCAAACAGCAGAGGAGGAGGAACCCCAAGGCGTATACAATGCACTGCAGAAAGA
CAAGATGGCAGAAGCCTACAGTGAGATCGGCACAAAAGGCGAGAGCGGAGAGGCAAGGGGCACGATGCC
CTTTACCAGGACAGCCACTTCCAAGCAGTGCAGTTCGGGAACAGAAGAGAGAGAAGGTTCAGAACTCA
CAAGGACCCCTTGGTTAAGAGCCCGCCCAAGGTGAAAGCACCCAGCAGAGTAGCCAATCCTGTGCCAG
CGTCTTACGATCCCCACTCTGTGGAGTCCATGGCCACCCAGTAGCAGCTCCAGCTC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >MR224932 representing NM_001113394
 Red=Cloning site Green=Tags(s)

MKWKVSVLACILHVRFPGEAQSFGLLDPKLCYLLDGILFIYGVIIITALYLRAKFSRSAETAANLQDPNQ
 LYNELNLGRREEYDVLEKKRARDPEMGGKQRRRNPNQEGVYNALQDKMAEAYSEIGTKGERRRGKGDG
 LYQDSHFQAVQFGNRREREGSELTRTLGLRARPKGESTQQSSQSCASVFSIPTLWSPWPPSSSSQL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_001113394

ORF Size: 618 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001113394.2](#), [NP_001106865.1](#)

RefSeq Size: 2957 bp

RefSeq ORF: 621 bp


Locus ID: 12503

Cytogenetics: 1 73.14 cM

MW: 23.3 kDa

Gene Summary: Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways. CD3Z ITAMs phosphorylation creates multiple docking sites for the protein kinase ZAP70 leading to ZAP70 phosphorylation and its conversion into a catalytically active enzyme. Plays an important role in intrathymic T-cell differentiation. Additionally, participates in the activity-dependent synapse formation of retinal ganglion cells (RGCs) in both the retina and dorsal lateral geniculate nucleus (dLGN). [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR224932