

Product datasheet for MR225016L3

Rfx3 (NM_001166414) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Rfx3 (NM_001166414) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Rfx3
Synonyms:	C230093O12Rik; MRFX3
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR225016).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.

ACCN:	NM_001166414
ORF Size:	2250 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001166414.1
RefSeq Size:	9146 bp
RefSeq ORF:	2250 bp
Locus ID:	19726
UniProt ID:	P48381
Cytogenetics:	19 C1
Gene Summary:	Transcription factor required for ciliogenesis and islet cell differentiation during endocrine pancreas development. Essential for the differentiation of nodal monocilia and left-right asymmetry specification during embryogenesis. Required for the biogenesis of motile cilia by governing growth and beating efficiency of motile cells (PubMed:15121860, PubMed:19671664). Also required for ciliated ependymal cell differentiation (PubMed:16930429). Together with RFX6, participates in the differentiation of 4 of the 5 islet cell types during pancreas development, with the exception of pancreatic PP (polypeptide-producing) cells (PubMed:17229940). Regulates transcription by forming a heterodimer with another RFX protein and binding to the X-box in the promoter of target genes (By similarity). Regulates the expression of genes involved in ciliary assembly (DYNC2LI1, FOXJ1 and BBS4) and genes involved in ciliary motility (DNAH11, DNAH9 and DNAH5). Represses transcription of MAP1A in non-neuronal cells but not in neuronal cells. [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR225016L3