

Product datasheet for MR225061L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Eri1 (NM 026067) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Eri1 (NM_026067) Mouse Tagged ORF Clone Lentiviral Particle

Symbol:

3'hexo; 3110010F15Rik; eri-1; Thex1 Synonyms:

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Myc-DDK Tag: NM 026067 ACCN:

ORF Size: 1035 bp

ORF Nucleotide

OTI Disclaimer:

Cytogenetics:

Sequence:

The ORF insert of this clone is exactly the same as(MR225061).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 026067.3, NP 080343.4

8 A4

RefSeq Size: 5067 bp RefSeq ORF: 1038 bp Locus ID: 67276 **UniProt ID:** Q7TMF2





Gene Summary:

RNA exonuclease that binds to the 3'-end of histone mRNAs and degrades them, suggesting that it plays an essential role in histone mRNA decay after replication. A 2' and 3'-hydroxyl groups at the last nucleotide of the histone 3'-end is required for efficient degradation of RNA substrates. Also able to degrade the 3'-overhangs of short interfering RNAs (siRNAs) in vitro, suggesting a possible role as regulator of RNA interference (RNAi). Binds with high affinity to the 3' side of the stem-loop structure and to the downstream cleavage product (DCP) of histone pre-mRNAs. Requires for binding the 5'-ACCCA-3' sequence present in stem-loop structure. Able to bind other mRNAs (By similarity). Required for 5.8S rRNA 3'-end processing. Also binds to 5.8s ribosomal RNA.[UniProtKB/Swiss-Prot Function]