

Product datasheet for MR225083L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Fnbp1 (NM_001038700) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Fnbp1 (NM 001038700) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Fnbp1

Synonyms: 1110057E06Rik; 2210010H06Rik; FBP1; Fbp17

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001038700

ORF Size: 1650 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR225083).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001038700.2, NP 001033789.1

RefSeq Size: 4545 bp
RefSeq ORF: 1653 bp
Locus ID: 14269
UniProt ID: Q80TY0

Cytogenetics: 2 B







Gene Summary:

Required to coordinate membrane tubulation with reorganization of the actin cytoskeleton during the late stage of clathrin-mediated endocytosis. Binds to lipids such as phosphatidylinositol 4,5-bisphosphate and phosphatidylserine and promotes membrane invagination and the formation of tubules. Also enhances actin polymerization via the recruitment of WASL/N-WASP, which in turn activates the Arp2/3 complex. Actin polymerization may promote the fission of membrane tubules to form endocytic vesicles. May act as a link between RND2 signaling and regulation of the actin cytoskeleton. May be required for the lysosomal retention of FASLG/FASL (By similarity).[UniProtKB/Swiss-Prot Function]