

Product datasheet for MR225090

Taar1 (NM_053205) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Taar1 (NM_053205) Mouse Tagged ORF Clone

Tag: Myc-DDK

Symbol: Taar1

Synonyms: taR-1; Tar1; Trar1

Mammalian Cell Neomycin

Selection:

Vector: pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

ORF Nucleotide >MR225090 representing NM_053205

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Sequence: >MR225090 representing NM_053205

Red=Cloning site Green=Tags(s)

MHLCHAITNISHRNSDWSREVQASLYSLMSLIILATLVGNLIVIISISHFKQLHTPTNWLLHSMAIVDFL LGCLIMPCSMVRTVERCWYFGEILCKVHTSTDIMLSSASIFHLAFISIDRYCAVCDPLRYKAKINISTIL VMILVSWSLPAVYAFGMIFLELNLKGVEELYRSQVSDLGGCSPFFSKVSGVLAFMTSFYIPGSVMLFVYY RIYFIAKGQARSINRTNVQVGLEGKSQAPQSKETKAAKTLGIMVGVFLVCWCPFFLCTVLDPFLGYVIPP SLNDALYWFGYLNSALNPMVYAFFYPWFRRALKMVLLGKIFQKDSSRSKLFL


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mm9020 e06.zip

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_053205

ORF Size: 996 bp

OTI Disclaimer:

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore,

OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts

OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at customercare te

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

ORÏGENE

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 053205.1</u>, <u>NP 444435.1</u>

38.1 kDa

 RefSeq Size:
 999 bp

 RefSeq ORF:
 999 bp

 Locus ID:
 111174

 UniProt ID:
 Q923Y8

 Cytogenetics:
 10 A4

MW:

Gene Summary: Receptor for trace amines, including beta-phenylethylamine (b-PEA), p-tyramine (p-TYR),

octopamine and tryptamine, with highest affinity for b-PEA and p-TYR. Unresponsive to classical biogenic amines, such as epinephrine and histamine and only partially activated by dopamine and serotonin. Trace amines are biogenic amines present in very low levels in

mammalian tissues. Although some trace amines have clearly defined roles as neurotransmitters in invertebrates, the extent to which they function as true


neurotransmitters in vertebrates has remained speculative. Trace amines are likely to be involved in a variety of physiological functions that have yet to be fully understood. The signal

transduced by this receptor is mediated by the G(s)-class of G-proteins which activate

adenylate cyclase.[UniProtKB/Swiss-Prot Function]



Product images:



Circular map for MR225090