

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR225256L4V

Neurod2 (NM_010895) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Neurod2 (NM_010895) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Neurod2
Synonyms:	bHLHa1; Ndrf
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_010895
ORF Size:	1149 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR225256).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 010895.3, NP 035025.3</u>
RefSeq Size:	3137 bp
RefSeq ORF:	1152 bp
Locus ID:	18013
UniProt ID:	<u>Q62414</u>
Cytogenetics:	11 61.75 cM



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Neurod2 (NM_010895) Mouse Tagged ORF Clone Lentiviral Particle - MR225256L4V

Gene Summary:Transcriptional regulator implicated in neuronal determination. Mediates calcium-dependent
transcription activation by binding to E box-containing promoter. Critical factor essential for
the repression of the genetic program for neuronal differentiation; prevents the formation of
synaptic vesicle clustering at active zone to the presynaptic membrane in postmitotic
neurons. Induces transcription of ZEB1, which in turn represses neuronal differentiation by
down-regulating REST expression. Plays a role in the establishment and maturation of
thalamocortical connections; involved in the segregation of thalamic afferents into distinct
barrel domains within layer VI of the somatosensory cortex. Involved in the development of
the cerebellar and hippocampal granular neurons, neurons in the basolateral nucleus of
amygdala and the hypothalamic-pituitary axis. Associates with chromatin to the DPYSL3 E
box-containing promoter.[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US