

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR225292L3V

Hif1an (NM_176958) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Hif1an (NM_176958) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Hif1an
Synonyms:	2310046M24Rik; A830014H24Rik; FIH; FIH1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_176958
ORF Size:	1047 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR225292).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 176958.3, NP 795932.2</u>
RefSeq Size:	6190 bp
RefSeq ORF:	1050 bp
Locus ID:	319594
UniProt ID:	<u>Q8BLR9</u>
Cytogenetics:	19 C3



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Hydroxylates HIF-1 alpha at 'Asn-799' in the C-terminal transactivation domain (CAD).Functions as an oxygen sensor and, under normoxic conditions, the hydroxylation prevents
interaction of HIF-1 with transcriptional coactivators including Cbp/p300-interacting
transactivator. Involved in transcriptional repression through interaction with HIF1A, VHL and
histone deacetylases. Hydroxylates specific Asn residues within ankyrin repeat domains (ARD)
of NFKB1, NFKBIA, NOTCH1, ASB4, PPP1R12A and several other ARD-containing proteins. Also
hydroxylates Asp and His residues within ARDs of ANK1 and TNKS2, respectively. Negatively
regulates NOTCH1 activity, accelerating myogenic differentiation (By similarity). Positively
regulates ASB4 activity, promoting vascular differentiation.[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US