

# Product datasheet for MR225313L3V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Flt4 (NM\_008029) Mouse Tagged ORF Clone Lentiviral Particle

#### **Product data:**

Product Type: Lentiviral Particles

**Product Name:** Flt4 (NM\_008029) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Flt4

Synonyms: Al323512; Chy; Flt-4; VEGFR-3; VEGFR3

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 008029

ORF Size: 4089 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(MR225313).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 008029.3, NP 032055.1

 RefSeq Size:
 5853 bp

 RefSeq ORF:
 4092 bp

 Locus ID:
 14257

 UniProt ID:
 P35917

Cytogenetics: 11 29.69 cM







#### **Gene Summary:**

Tyrosine-protein kinase that acts as a cell-surface receptor for VEGFC and VEGFD, and plays an essential role in adult lymphangiogenesis and in the development of the vascular network and the cardiovascular system during embryonic development. Promotes proliferation, survival and migration of endothelial cells, and regulates angiogenic sprouting. Signaling by activated FLT4 leads to enhanced production of VEGFC, and to a lesser degree VEGFA, thereby creating a positive feedback loop that enhances FLT4 signaling. Modulates KDR signaling by forming heterodimers. Mediates activation of the MAPK1/ERK2, MAPK3/ERK1 signaling pathway, of MAPK8 and the JUN signaling pathway, and of the AKT1 signaling pathway. Phosphorylates SHC1. Mediates phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase. Promotes phosphorylation of MAPK8 at 'Thr-183' and 'Tyr-185', and of AKT1 at 'Ser-473'.[UniProtKB/Swiss-Prot Function]