

## Product datasheet for **MR225318**

### Rab3a (NM\_001166399) Mouse Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Rab3a (NM\_001166399) Mouse Tagged ORF Clone  
**Tag:** Myc-DDK  
**Symbol:** Rab3a  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**ORF Nucleotide Sequence:** >MR225318 ORF sequence  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGATCGCC**

ATGGCTTCGCCACAGACTCTCGCTATGGGCAGAAGGAGTCTCAGACCAGAACTTCGACTATATGTTCA  
 AGATCCTGATCATTGGGAACAGCAGCGTGGGCAAAACCTCGTTCCTCTCCGCTACGCAGATGACTCCTT  
 CACTCCAGCCTTTGTGACACCGTTGGCATAGACTTCAAGGTCAAACCATCTACCGCAACGACAAGAGG  
 ATCAAGCTGCAGATCTGGGACACAGCAGGGCAAGAGCGGTACCGCACCATCACCACAGCCTATTACCGAG  
 GCGCCATGGGCTTCATCCTAATGTATGACATCACCAATGAGGAGTCATTTAATGCAGTGCAGGACTGGTC  
 CACTCAGATCAAAACTTACTCGTGGGACAATGCCAGGTGCTGCTGGTGGGAAACAAGTGTGACATGGAA  
 GATGAGCGAGTGGTGTCTCAGAGCGTGGCCGGCAGCTGGCTGACCACCTGGGCTTTGAGTCTTTGAGG  
 CCAGCGCCAAGGACAACATTAATGTCAAGCAGACGTTTGAACGTCTGGTGGACGTGATCTGTGAGAAGAT  
 GTCAGAGTCCCTGGATACTGCAGACCCTGCGGTACCGGTGCCAAGCAGGGCCCGCAGCTCACCACCGAG  
 CAGGCGCCACCTCATCAGGATTGTGCCTGC

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >MR225318 protein sequence  
 Red=Cloning site Green=Tags(s)


MASATDSRYGQKESSDQNFDMFKILIIIGNSSVGKTSFLFRYADDSFTPAFVSTVGIDFKVKTIYRNDKR  
 IKLQIWDTAGQERYRTITTAYYRGAMGFILMYDITNEESFNAVQDWSTQIKTYSWDNAQVLLVGNKCDME  
 DERVVSSERGRQLADHLGFEFFEASAKDNINVKQTFERLVDVICEKMSESLDTADPAVTGAKQGPQLTDQ  
 QAPPHQDCAC

**TR**TRPLEQKLISEEDLAANDILDYKDDDDKV


[View online »](#)

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**



**ACCN:** NM\_001166399

**ORF Size:** 660 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

**RefSeq:** [NM\\_001166399.3](#)


**RefSeq Size:** 1490 bp

**RefSeq ORF:** 663 bp

**Locus ID:** 19339  
**UniProt ID:** [P63011](#)  
**Cytogenetics:** 8 34.15 cM  
**MW:** 25 kDa

**Gene Summary:** Small GTP-binding protein that plays a central role in regulated exocytosis and secretion. Controls the recruitment, tethering and docking of secretory vesicles to the plasma membrane (PubMed:11598194). Upon stimulation, switches to its active GTP-bound form, cycles to vesicles and recruits effectors such as RIMS1, RIMS2, Rabphilin-3A/RPH3A, RPH3AL or SYTL4 to help the docking of vesicles onto the plasma membrane (By similarity). Upon GTP hydrolysis by GTPase-activating protein, dissociates from the vesicle membrane allowing the exocytosis to proceed (By similarity). Stimulates insulin secretion through interaction with RIMS2 isoform RIMS2 and RPH3AL effectors in pancreatic beta cells (PubMed:15159548, PubMed:20674857). Regulates calcium-dependent lysosome exocytosis and plasma membrane repair (PMR) via the interaction with 2 effectors, SYTL4 and myosin-9/MYH9 (By similarity). Acts as a positive regulator of acrosome content secretion in sperm cells by interacting with RIMS1 (By similarity). Plays a role in the regulation of dopamine release by interacting with synaptotagmin I/SYT (By similarity).[UniProtKB/Swiss-Prot Function]

## Product images:



Circular map for MR225318