

## Product datasheet for **MR225337L3V**

### **Pkd2l1 (NM\_181422) Mouse Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Pkd2l1 (NM_181422) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Pkd2l1
Synonyms:	B830002B15; BC046386; PCL; PKD2L; PkdI; TRPP3
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_181422
ORF Size:	2280 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR225337).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_181422.3</a> , <a href="#">NP_852087.2</a>
RefSeq Size:	3321 bp
RefSeq ORF:	2283 bp
Locus ID:	329064
UniProt ID:	<a href="#">A2A259</a>
Cytogenetics:	19 36.91 cM



[View online »](#)

**Gene Summary:**

Pore-forming subunit of a heteromeric, non-selective cation channel that is permeable to Ca(2+) (PubMed:16891422, PubMed:15548533, PubMed:19464260, PubMed:20538909, PubMed:21185261, PubMed:22420714, PubMed:25820328, PubMed:28904867, PubMed:29567962). Pore-forming subunit of a calcium-permeant ion channel formed by PKD1L2 and PKD1L1 in primary cilia, where it controls cilium calcium concentration, but does not affect cytoplasmic calcium concentration (PubMed:24336288, PubMed:24336289). The channel formed by PKD1L2 and PKD1L1 in primary cilia regulates sonic hedgehog/SHH signaling and GLI2 transcription (PubMed:24336288). Pore-forming subunit of a channel formed by PKD1L2 and PKD1L3 that contributes to sour taste perception in gustatory cells (PubMed:16891422, PubMed:16929298, PubMed:20406802, PubMed:21098668, PubMed:21625513). The heteromeric channel formed by PKD1L2 and PKD1L3 is activated by low pH, but opens only when the extracellular pH rises again (PubMed:18535624, PubMed:19464260, PubMed:20538909, PubMed:20406802, PubMed:22420714, PubMed:28904867, PubMed:29567962). May play a role in the perception of carbonation taste (PubMed:19833970). May play a role in the sensory perception of water, via a mechanism that activates the channel in response to dilution of salivary bicarbonate and changes in salivary pH (PubMed:28553944).[UniProtKB/Swiss-Prot Function]