

## Product datasheet for MR225368L3

### Pou4f1 (NM\_011143) Mouse Tagged Lenti ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Pou4f1 (NM_011143) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Pou4f1
Synonyms:	Brn-3; Brn-3.0; Brn3; Brn3.0; Brn3a; E130119J07Rik
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR225368).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.

ACCN:	NM_011143
ORF Size:	1263 bp




[View online »](#)

<b>OTI Disclaimer:</b>	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:custsupport@origene.com">custsupport@origene.com</a> or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a></p>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>RefSeq:</b>	<a href="#">NM_011143.4</a> , <a href="#">NP_035273.3</a>
<b>RefSeq Size:</b>	3801 bp
<b>RefSeq ORF:</b>	1266 bp
<b>Locus ID:</b>	18996
<b>UniProt ID:</b>	<a href="#">P17208</a>
<b>Cytogenetics:</b>	14 E2.3

**Gene Summary:**

Multifunctional transcription factor with different regions mediating its different effects (PubMed:10640682, PubMed:8621561, PubMed:9694219, PubMed:9722627). Acts by binding (via its C-terminal domain) to sequences related to the consensus octamer motif 5'-ATGCAAAT-3' in the regulatory regions of its target genes (PubMed:8621561, PubMed:17668438). Regulates the expression of specific genes involved in differentiation and survival within a subset of neuronal lineages. It has been shown that activation of some of these genes requires its N-terminal domain, maybe through a neuronal-specific cofactor (PubMed:12934100). Activates BCL2 expression and protects neuronal cells from apoptosis (via the N-terminal domain) (PubMed:9722627). Induces neuronal process outgrowth and the coordinate expression of genes encoding synaptic proteins (PubMed:8972215). Exerts its major developmental effects in somatosensory neurons and in brainstem nuclei involved in motor control. Stimulates the binding affinity of the nuclear estrogen receptor ESR1 to DNA estrogen response element (ERE), and hence modulates ESR1-induced transcriptional activity (PubMed:9448000). May positively regulate POU4F2 and POU4F3 (PubMed:8876243). Regulates dorsal root ganglion sensory neuron specification and axonal projection into the spinal cord (PubMed:22326227). Plays a role in TNFSF11-mediated terminal osteoclast differentiation (PubMed:17668438). Negatively regulates its own expression interacting directly with a highly conserved autoregulatory domain surrounding the transcription initiation site (PubMed:12441296).[UniProtKB/Swiss-Prot Function]

**Product images:**


Circular map for MR225368L3